MEMBRANE ULTRASTRUCTURE IN NERVE CELLS

H. Fernández-Morán, M.D., Ph.D.

Intensive Study Program in the Neurosciences

Boulder, Colorado -- Summer 1966

For the Neurosciences Research Program

Massachusetts Institute of Technology

The Rockefeller University Press



Department of Biophysics
The University of Chicago
Chicago, Illinois

MEMBRANE ULTRASTRUCTURE IN NERVE CELLS

by Humberto Fernandez-Moran, M.D., Ph.D.

Department of Biophysics, University of Chicago Chicago, Illinois

appearing in
The Neurosciences, A Survey
for Synthesis,
based on the NRP Summer, 1966
Intensive Study Program in
the Neurosciences,
to be published by the
Rockefeller University Press,
New York

Introduction

Membranes and their derivatives play a key role in all biological systems, constituting about 40% to 90% of the total mass in different cell types. They represent a major structural component of nervous tissues, ranging from individual limiting membranes of the wide variety of cells to multiply folded or invaginated systems which are often disposed in "paracrystalline" arrays. 1,2,3 Attention has focused on these specialized lamellar systems, including mitochondria and photoreceptors, as prototypes of organized membranes which carry out energy transduction function with remarkable efficiency. The highly ordered structural framework of these cell membranes provides relatively stable "modular floor-space," thereby ensuring periodic arrangement of specialized macromolecular enzyme complexes and related assemblies. 4,5,6

It is possible, in light of accumulating evidence, that these membrane systems may operate equally well as information transducers. In fact, the neuron, which as the unit of the nervous system is invested with the primary function of information transfer, possesses membrane-bound terminal endings with specialized synaptic structures that functionally interlink specific cell groups. The suggested location of subcellular or molecular switch gears at such synaptic junctions manifests their integral association

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Morán with the processes of information storage, transfer, and readout. According to Schmitt 7 the primary role of molecular recognition of coded information stored in macromolecules of the brain cells, is the formation of the exceedingly complex neuronal network of the human brain. Some of the most sophisticated behavioral patterns of the instinctual type involve primarily molecular recognition of coded information stored in DNA-RNA-protein macromolecular systems. The underlying molecular specificity in the key junctional regions of the nervous system must ultimately be taken into account in any attempts to explain the singular capacity for accurate storage of the near infinite number of memories which span a lifetime.

The pioneering polarized light 8 and x-ray diffraction investigations of the nerve myelin sheath 9 provided a standard of reference for the study of cell membrane ultrastructure in general. However, the task of directly visualizing individual cell membranes, whose thickness is in the order of 100 Å to 200 Å, remained for the higher resolving power of the electron microscope. 10,11 Ultrastructural analysis is now progressing beyond the initial stage of delineating the membrane's lipoprotein framework.

Based essentially on the paucimolecular theory of membrane structure proposed by Danielli and Davson, ¹² Robertson has propounded a unit membrane hypothesis. ¹³ After demonstrating that plasma membranes of all cells and the membranes of

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Morán their organelles show certain common structural features, he proposes that biological membranes are comprised of a phospholipid bilayer system, 40 to 60 Å thick, which

is coated on both sides with monolayers of macromolecular

material, presumed to be proteins.

This unit membrane concept of Robertson serves as a useful working hypothesis which must now be extended and modified according to the large body of evidence provided by correlated electron microscopic and biochemical investigations during the past few years. These studies have revealed specific macromolecular functional repeating units and the counterpart structural elements in mitochondria, chloro-19,20 plasts, retinal rod outer segments, and in a large 24,25,26,27. number of membranes from different cell types.

Investigations of nerve membranes and related structures were recently carried out by high resolution electron microscopy, using improved preparation and instrumentation techniques, which have already proved successful in the analysis of virus fine structure and other biological systems. Increasing emphasis has been placed on the study of the detailed organization of the enzyme and multienzyme complexes intimately associated with cell membranes as the molecular componentry ultimately responsible for the highly specific energy and information transduction functions.

Membrane Ultrastructure in Nerve Cells - H. Fernández-Moran

Only salient aspects can be given here of the available evidence bearing on organization within the plane of the layers, and the existence of macromolecular repeating units in membranes. Particular attention will be paid to correlated biochemical and electron microscopic studies of well characterized multienzyme complexes which are representative of the types of macromolecular assemblies that may be found in association with nerve membranes. An attempt will be made to show how this association of specific macromolecular assemblies with the hydrated lipoprotein framework of the nerve membranes may serve to fulfill many structural and functional requirements which are unique to the nervous system. Representative examples will illustrate the organization of the nerve myelin sheath and junctional regions of the nervous system. The role of water as an integral structural component will also be discussed in relation to the functional specificity of these key sites of the nervous system.

Organization of Myelin Membranes

Elucidation of the general submicroscopic organization of the nerve myelin sheath may be regarded as the corner stone of membrane ultrastructure research. Nearly two decades ago, F. O. Schmitt and his associates^{9,28} gave the first detailed account of the highly ordered, layered structure of the myelin sheath, as deduced from their classic small-angle, x-ray⁹ diffraction studies of living nerve and earlier polarized

Membrane Ultrastructure in Nerve Cells - H. Fernández-Moran

light analysis. The postulated regular, concentric arrangement of the myelin sheath layers was subsequently verified by direct observation in the electron microscope. 10,11,14,30 Geren and Schmitt then showed that the myelin sheath derives from invaginated Schwann cell membranes multiply wrapped around the axon of embryonic peripheral nerve fibers. In the morphogenetic process of myelination, leading to the formation of this condensed, "liquid crystalline" membrane system, the constituent lipid and protein components in the limiting membranes of both axons and Schwann cells (or glia in the central nervous system) play specific roles.

The x-ray diffraction pattern of fresh intact nerve, which serves as a standard of reference for the study of myelin structure, features a number of very well-defined, low-angle reflections. These reflections exhibit a characteristic alternation of intensities in the even and odd orders, and can be accounted for as the first five orders of a fundamental spacing, varying from about 170 Å in amphibians to 178 Å (Fig. 1A) to 184 Å in mammalian peripheral nerve. The low-angle x-ray diffraction patterns recorded from normal nerve furnish general information on the dimensions and approximate distribution of scattering groups in the radial direction of the myelin sheath. Earlier studies of physical and chemical modifications in the myelin sheath established a correspondence between the layer spacings observed in the electron micrographs (Fig. 1A) and the fundamental radial

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran repeating unit indicated by the low-angle x-ray diffraction patterns (Fig. 1c)

It should be pointed out, however, that the concentric laminated structure observed in electron micrographs of thin nerve sections is simply a pattern of the selective deposition of osmium or other reagents at certain sites, and it cannot be interpreted in terms of specific regions containing lipids, lipoproteins, or protein constituents of the myelin sheath. Moreover, extensive modifications are introduced by complex fixation, dehydration and embedding artifacts which must be taken into account in a critical evaluation of the fine structures observed in thin sections. 33,34 With these reservations in mind, it is assumed that the framework of the fundamental radial unit of the sheath is formed by two adjacent bimolecular layers of mixed lipids about 60 A in thickness; each bilayer is coated with sheets of hydrated protein or other polysaccharide, non-lipid macromolecular material. The two bimolecular layers are distinguished by a "difference factor" which can be accounted for by assuming that in the process of myelin formation, the asymmetric wrapping of the Schwann cell membrane around the axon will produce a symmetry difference in successive layers.

Lipid Composition of Myelin

Relatively little is known about the composition of the major lipids and their fatty acids in myelin, since this component is very difficult to isolate reliably and in sufficient

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran quantity. Recently, however, J. S. O'Brien and associates 35 have reported the results of their efforts to determine the composition of lipids and fatty acids obtained from carefully isolated myelin, gray matter, and white matter in normal human frontal lobes at various ages. The lipid content of myelin, ranging between 78% and 81% of the dry weight, was found to be much higher than the lipid content of gray or white matter. According to the pioneering studies of Folch-Pi and other earlier workers, myelin contains much higher molar proportions of cerebroside sulfate and plasmalogenes and slightly higher molar proportions of cholesterol than gray matter. It is noteworthy that in myelin the content of sphingolipids, containing very longchain fatty acids (19 to 26 carbon atoms), is proportionately ten times that of any other membrane structure analyzed. Also, 1 in 17 fatty acids is polyunsaturated in the myelin group of lipids, while in the gray matter group the corresponding value is 1 in 5.

The recent molecular model of myelin proposed by Vandenheuvel ³⁷ serves to illustrate certain features characteristic of myelin lipids which could account for the highly stable membrane structure. Lipids in myelin are not covalently bound. The three major forces holding these molecules within

the membrane result from: 1) electrostatic interactions between polar groups of lipids and oppositely charged groups in adjacent proteins, 2) hydrogen bonding between oxygen and nitrogen atoms in lipids and adjacent proteins, and 3) London-van der Waals dispersion forces between CH, pairs in hydrocarbon tails of adjacent lipid molecules. latter may be the major force holding such molecules together, since the total force owing to the additive interactions of the large number of CH, pairs is appreciable. On the basis of steric hindrance, it can be expected that lipids containing polyunsaturated acids will not be held as rigidly in a bimolecular leaflet, thus leading to a more free and less-stable structure. The higher proportion of saturated glycerophosphatides present in myelin will therefore impart greater stability, since the lipid molecules can be more closely packed than in the membrane structures of gray matter which have a significantly higher content of polyunsaturated glycerophosphatides. Finally, the sphingolipids containing long-chain fatty acids of the type found in myelin can form interdigitated cholesterol-sphingo-myelin complexes as proposed by Vandenheuvel. All of this new data makes it possible to account for the closely packed and highly ordered myelin membrane structure. It may also help explain the remarkable degree of metabolic inertness of which has been derived from turnover myelin lipids 38,39 studies with labelled lipids. Myelin is thus considered to

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran be the most stable membrane known, lasting the lifetime of the animal.

All of these conclusions relate mainly to the lipid composition, but our knowledge about the structure, composition, and function of other constituents of myelin, including proteins, water, polysaccharides and salts is still very meager. Recent experiments by M. Singer and M. Salpeter indicate that tritium-labelled L-histidine is transported through the Schwann and myelin sheath into the axon of peripheral nerves. Further studies along these lines should help clarify how incorporated amino acids traverse the myelin sheath.

Schmitt has emphasized that the unique properties of the axon surface membrane appear to be determined by the inner protein component which is considered to be different from the bimolecular lipid phase and the external non-lipid layer. The inner monolayer of globular protein molecules, forming a continuous, mosaic sheet, may be capable of responding to electrical and ionic changes with fast conformation changes of the protein molecules. Such dynamic changes of protein molecules in relation to the adjoining lipid bilayer may significantly alter the net fixed charge, and consequently, the membrane or action potential. Schmitt and co-workers have given the name "electrogenic protein" to this membrane-associated protein capable of rapid conformation change in response to electric field or other bioelectric

Membrane Ultrastructure in Nerve Cells - H. Fernández-Morán modifications. Identification and localization of the electrogenic protein may eventually be possible by combining the immuno-neurological experiments of Huneeus-Cox et al 42 with appropriately labelled antibody techniques for electron microscopy, possibly using cytochrome-C which is smaller than ferritin. The typical combination of the antibody or -SH reagent with the electrogenic protein may also make it possible to introduce selective labelling with heavy metals for high resolution electron microscopy.

Subunit Structure of Myelin Membranes

Earlier investigations 10,11,14,30,33 had shown that the layers of the myelin sheath tend to dissociate into granular or rod-shaped particles about 60 A. This transformation occurs following a wide variety of controlled physical and chemical treatments which include enzymatic digestion, freezing and thawing of nerve, and low-temperature fixation and embedding techniques (Fig. 1A). Correlated electron microscopic and x-ray diffraction data also indicated the possibility of a regular organization within the plane of the layers, probably involving units of 60 to 80 Å. 33 In vitro nerve degeneration studies 4,6 have disclosed various forms of granular dissociation in the lamellae which closely resemble structures observed after enzymatic digestion. It was determined from low-angle, x-ray diffraction patterns that modifications of the x-ray scattering power within the radial unit corresponded with the rate of degeneration and were likewise consistent with the presence of a granular structure in the

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran dense and intermediate layers of the myelin sheath. These results and consistent electron microscopic findings indicate a regular fine structure within the concentric arrays of the dense and intermediate layers (Fig. 1A) which warrants careful consideration.

Our investigations made use of cryofixation techniques which are based on rapid freezing of fresh or glycerinated specimens with liquid helium II at 1 to 20 K, or with under-4,6 cooled nitrogen. This is followed by freeze-substitution and embedding in plastics at low temperatures. Conditions are maintained which minimize ice-crystal formation and artifacts of chemical fixation, extraction, and embedding. These and related techniques for thin sectioning have consistently yielded better morphological and histochemical preservation of all types of lamellar systems and membrane derivatives than the standard procedures. As shown in Fig. 1A, particulate subunit structures of 60 to 80 Å are regularly found, mainly within the plane of the intermediate layers, but also in the dense layers of the wellpreserved concentric laminated structure. This type of structure was seen informally in all kinds of peripheral and central optic nerves, particularly of frog and rats. It is detected in cross sections, longitudinal sections, and particularly in tangential sections revealing face views of the plane of the layers.

Special techniques were also applied in preparing ultrathin, frozen sections of fresh tissue. They were cut with a special Page - 11

Membrane Ultrastructure in Nerve Cells - H. Fernández-Morán microtome and diamond knife in a cryostat at -30 to -180° C. The specimens were often examined directly without thawing by embedding in vitrified heavy metal layers. A liquid nitrogen cooling device was used in conjunction with appropriate low-temperature electron microscopy techniques. Ultrathin frozen sections of fresh frog sciatic nerve, negatively stained with phosphotungstate (Fig. 2B), show characteristic repeating particulate units (arrows) of about 50 to 60 Å. These units, mainly localized in the intermediate layers and attached to the dense myelin layers, were consistently found in the negatively stained, unfixed specimens. Bearing in mind inherent artifact possibilities (e.g., myelin figures), these and related methods are now being further investigated in an attempt to examine the fine structure of the myelin sheath under conditions closely approximating its native hydrated state.

Recent work by Branton and Moor 45 using the freeze-etching technique show that the fracture faces of completely myelinated membranes generally appear smooth and relatively free of particles, in contrast to the inner faces of all other membranes which show distinct particulate components. These findings are of considerable significance because no heavy metal staining is involved. However, pretreatment of nerves in 20% glycerol, followed by freezing, is seen in our earlier low-temperature, x-ray diffraction studies 4,6 to produce characteristic modifications in the hydrated lipo-

Membrane Ultrastructure in Nerve Cells - H. Fernández-Morán protein constituents, often with underlying drastic alterations of the hydrated matrix permeating the entire sheath. In view of the limited level of resolution and the degree of possible artifacts inherent in freeze etching when dealing with such fine structures, further work is clearly indicated.

Organization of Membranes in Photoreceptors

Membranes of the vertebrate photoreceptors comprise "unit discs" which are stacked in regular order in the outer segments. Earlier polarized light studies indicated that the rod outer segments consist of thin, transversely arranged protein layers which alternate with longitudinally oriented layers of lipid molecules. Electron microscopy has confirmed this idea by revealing the fine structure of the outer segment and its composition of several hundred unit discs about 150 to 200 % thick. 2,4,30,49

Improved low-temperature electron microscopy techniques revealed an intermediate layer between the dense layers, and electron-dense structures within the layers which appear as globular subunits (Fig. 3A).^{6,21} These subunits look well organized in dark-adapted outer segments and may be related to the presence of the photopigment complexes, as suggested by data from polarized light analysis and experimental modifications. Since rhodopsin represents about 40% of the dry weight in frog retinal rod outer segments, it may be regarded as a principal structural

Membrane Ultrastructure in Nerve Cells - H. Fernández-Morán component of the photoreceptor membrane system.

Because of its inherent uncertainties, however, electron microscopical data on the fine structure of membrane in vivo cannot be considered reliable unless confirmed by an independent technique such as x-ray diffraction analysis. Previous studies of this type 50 demonstrated a spacing of about 320 A which was tentatively correlated with the repeating units observed along the axis of the rod outer segments in osmium-fixed preparations. Although these lowangle, x-ray diffraction patterns revealed layering of the membranes and the intermembrane spacings, they did not yield any specific information on the structure within the plane of the membranes. Recent low-angle, x-ray diffraction studies by Blasie and Dewey, Blaurock and Worthington, 51 have substantiated the presence of an orderly, globular substructure within the isolated outer segment membranes of photoreceptors in frog retina.

The x-ray diffraction patterns gave reflections which are consistent with a square array of particles within the membranes with a unit cell size of 70 Å. Low-angle, x-ray diffraction showed periodicities in isolated outer segments to be identical with those of intact outer segments from the retinal receptors. The authors also observed an orderly arrangement of particles about 40 Å in diameter within negatively stained preparations of outer segment

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran membranes. However, our limited knowledge of the chemical composition of photoreceptors precludes interpretation of these particulate subunits in terms of specific photopigments, or associated multienzyme components.

We have used special preparation techniques and low-temperature electron microscopy to examine outer segments of the frog retinal rod. Fresh, ultrathin sections of light and dark-adapted segments were prepared without fixing and without embedding. The specimens were kept frozen or dried and were stained, negativelyor positively, with buffered phosphotungstate, uranyl formate, and other electron-dense reagents. Additional structural detail can be detected in these well-preserved preparations (Fig. 3B,C). The globular subunits (60 to 70 Å), which are prominent in the dense and intermediate layers, appear to be built up of minute particles, about 10 to 15 Å, regularly arranged in clusters or rows. Symmetrical alignment of these dense particles is usually maintained, even in certain regions where the dense layers are split.

These preparation techniques, which combine the advantages of thin sectioning with the enhanced preservation of negative staining applied to native, unextracted specimens, 6,15 are particularly suitable for correlated biochemical studies.

Moreover, when used in combination with enzymological assays of isolated retinal outer segments it should be possible

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Morán to obtain essential data on the localization of the visual pigments, with associated lipoprotein components and specific enzyme systems.

Invertebrate Photoreceptors

Invertebrate photoreceptors are essentially composed of highly ordered tubular compartments which are considered differentiated membrane extensions, or microvilli, of the retinula cell membrane. Closely packed arrays of these tubules (ca. 400 to 1,000 Å in diameter) comprise the radially disposed rhabdomeres of the retinula cells in each ommatidium of the insect compound eye.

Earlier electron microscope studies and more recent electrophysiological investigations by Waterman et al⁵⁶ point to the complex arrangement of photopigments within the plane of the membranes as a key to an integral understanding of the photoreceptor system. For example, the insect eye's remarkable capacity for analyzing polarized light depends essentially on the dichroism of the photopigment molecules, which must possess a highly ordered "paracrystalline" arrangement within the microvilli in order to provide the molecular basis for differential sensitivity to linearly polarized light. The precise orientation of the many thousands of closely packed tubular compartments in a single rhabd-omere comprise a unit dichroic analyzer at the cellular and subcellular levels. Recent data by Waterman and Horch, ⁵⁶

Membrane Ultrastructure in Nerve Cells - H. Fernández-Morán which includes a model for a two-channel polarization analyzer, indicates that one ommatidium suffices for detecting small e-vector orientations.

The differentiated membrane-bound components in these photo-receptors display a highly ordered and symmetrical arrangement throughout the various dimensional levels of organization. Ultimately, however, this exquisitely built, compact system owes its capability of extracting optimum information, through selective interaction with incoming light signals, to the precise three-dimensional arrangement of the photopigment molecules within the membranes.

Macromolecular Repeating Units of Structure and Function in Mitochondrial Membranes

A large body of evidence has accumulated during the past decade 4,6,15,16,17,18 showing that the electron transfer system, the respiratory transformations, and certain other systems of the mitochondrion are built up from characteristic macromolecular complexes. Large scale isolation of stable mitochondria under conditions which do not impair their main enzymatic activities, play a significant role by making these membranous organelles and their constituents available for biochemical analysis.

Correlated electron microscopic and biochemical studies 6,15

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran

have resulted in the detection of a macromolecular repeating particle associated with the cristae and the inner membrane of the external mitochondrial envelope. Many thousands (ca. 10⁴ to 10⁵) of these particles, are found disposed in regular arrays in a single mitochondrion using negative staining and other preparation techniques. The repeating particle, designated the mitochondrial elementary particle (EP), consist of three parts: a spherical or polyhedral head piece, 80 to 100 Å in diameter; a stalk (ca. 50 Å long); and a base piece (40 x 110 A). As shown in Figures 4 A.B the base pieces of the elementary particles form an integral part of the outer dense layers of the cristae. These characteristic repeating particles are particularly well demonstrated in negatively stained specimens of mitochondria in situ, of isolated mitochondria, and of submitochondrial membrane fragments with a complete electron transfer chain.

Following the first observations by Fernandez-Moran⁶,14,15 numerous investigators¹⁶,17,18,60,61 confirmed the existence of the repeating particles and underlying subunit organization of mitochondrial membranes using a wide variety of techniques, including the improved freeze-etching method applied by Moor and Muhlethaler.⁶² Since this technique operates on native frozen specimens and does not involve the use of chemical fixatives or stains, the successful demonstration of globular repeating units in mitochondria and other membrane systems has effectively disposed of tentative suppositions⁶³ that these characteristic units might be preparative artifacts.

Membrane Ultrastructure in Nerve Cells - H. Fernández-Morán

64,65,66,61.

It is, therefore, now generally agreed by most workers that these mitochondrial subunits exist, although the observed structural details will depend on the preparation techniques and experimental conditions.

Detection of these structural subunits of mitochondrial membranes suggested the existence of counterpart macromolecular functional units. A particulate unit containing a complete electron transfer chain was isolated from beef heart mitochondria. 15,59 The repeating unit has a molecular weight of about 1.3 x 10⁶ (corresponding to approximately 40 molecules of protein and 400 molecules of phospholipids). The experimental evidence obtained from our studies was consistent with the notion that the isolated (Fig. 4C) and reconstituted particles could be identified with the elementary particles (ER) visualized in situ. It was also assumed that the isolated elementary particles contained the complete electron transfer chain.

Although subsequent studies by Green and Perdue 17 indicate that there may be multiple species of mitochondrial elementary particles, the basic concept of a macromolecular repeating unit of mitochondrial structure and function is now generally accepted. The elementary particle of the mitochondrion is believed to be a prototype of a class of functional particles or macromolecular assemblies found in association with membranes generally.

Membrane Ultrastructure in Nerve Cells - H. Fernández-Morán

Macromolecular Repeating Units in Chloroplast Membranes and

other Membranes

Investigations by Park and Pon and others 19,20 have led to the identification of repeating units in the chloroplast membranes which are called quantasomes and may be the morphological expression of the physiological photosynthetic unit. X-ray diffraction work by Kreutz also indicates that the chloroplast lamellae, are built up of repeating units about 71 X in diameter. The molecular weight of a single quantasome is 2×10^6 . The interesting relationship between the quantasome of chloroplasts and the elementary particle of mitochondria has been pointed out by Park, et al. 20 Both are associated with the electron-transport system in membranes and both contain certain cytochromes, substituted benzoquinones and non-heme iron. Morphological differences have been revealed, however, in that four or more subunits appear in the quantasome. Branton and Park show a second type of subunit, 110 Å in average diameter, which forms part of a matrix around the larger 175 A quantasome units. It is concluded that the chloroplast lamellae consist of a matrix within which are densely packed subunits that form the major constituents of the photosynthetic membrane.

Plasma membranes of liver cells, microvilli, microsomal membranes, 45 bacterial membranes, and synaptic membranes are among the different types in which repeating units have been observed. In general, there is evidence that such

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran macromolecular repeating units appear in all membranes and either make up the entire membrane or comprise a major part of the membrane framework in close association with the bilayer lipid structure. Although these repeating units are of variable shape and size, with a range of molecular weights from 50,000 to several million, they all have significant features in common with the macromolecular assemblies conceived by F. O. Schmitt as distinctive hierarchial units of structure and function (Fig. 4D) invested with unique properties.

It has been shown that enzymes constitute a large proportion of the total protein, particularly of specialized energy transducing mitochondrial membranes. This catalytic protein, may therefore be regarded as the key determinant of the structural and functional organization of membrane systems. In all membrane systems, including nerve membranes, specific enzymes or enzyme complexes are attached to, or otherwise intimately associated with, the basic membrane framework in repetitive patterns of exquisite, three-dimensional, stereospecific configuration.

The concept of macromolecular repeating units or assemblies in membranes can therefore, in many ways, be regarded as an extension of the unit membrane hypothesis of Davson, 12 13 Danielli, and Robertson, which represents a necessary and logical transition consistent with present biochemical and physico-chemical evidence.

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran

Macromolecular Organization of Enzyme Systems

A. Pyruvate and α-Ketoglutarate Dehydrogenation Complexes

A unique opportunity to correlate functional properties with ultrastructure has recently been provided by biochemical and electron microscopic studies of the pyruvate dehydrogenase 70 complex (PDC) of Escherichia coli. This well-characterized multienzyme complex, with a molecular weight of about 4.8 million, has been isolated in highly purified form and analyzed by Lester Reed and associates. 67,68

PDC catalyzes a multistage oxidative decarboxylation of pyruvate, ^{67,71} and is composed of three enzymes. Each molecule of the PDC complex contains about 16 molecules of pyruvate decarboxylase (M.W. 183,000), 8 molecules of dihydrolipoic dehydrogenase (M.W. 112,000), and about 64 molecules of lipoic reductase-transacetylase (LRT). Lester Reed and his associates have also succeeded in reconstituting the complex from the isolated enzymes. The concept of the organization of the PDC complex which emerged from these classical biochemical studies is that of an organized assembly of enzymes in which each of the constituent enzymes is specifically located to permit efficient implementation of a consecutive reaction sequence.

Our correlative electron microscope studies ^{69,70} have confirmed and extended this picture through direct visualization of the exquisite molecular architecture of the PDC complex and its components. The negatively stained PDC

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran complex appears in electron micrographs (Figure 5A) as polyhedrical particles of about 300 to 400 Å in diameter featuring a central tetrad (Figure 5B).

Electron microscopy of the isolated LRT aggregate revealed a typical tetrad structure which closely resembles the central tetrad of the native PDC complex and appears to determine its overall structure. According to the model derived from the biochemical and electron-microscopic data, lipoic reductase transacetylase (LTR) is composed of 64 identical subunits situated at the eight vertices of a cube, while the molecules of pyruvate decarboxylase and dihydrolipoic dehydrogenase are respectively aligned on the twelve edges 70,71 and in the six faces of the cube.

The macromolecular organization of the α -ketoglutarate dehydrogenase complex (KGDC) isolated from $\underline{\mathbf{E}}$. coli appears to be similar to that of the PDC. As shown by electron microscopy (Fig. 5C,D), the smaller polyhedrical particles, with a diameter of about 280 Å, feature a tetrad core surrounded by peripheral subunits, which are not clearly defined as in the PDC complexes. Recent studies by Reed and his associates indicate that the macromolecular

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran organization of the bacterial and mammalian PDC and KCDC multienzyme complexes are fundamentally similar, and may be governed by the principles of self-assembly which determine the functional organization of protein shells of regular viruses with icosahedral symmetry.

B. Organization of L-Glutamate Dehydrogenase

L-Glutamate dehydrogenase (GDH), which has a molecular weight of 1 million, consists of four major subunits. This molecule is of interest since it is known to be reversibly dissociated 74 by certain steroids and hormones such as thyroxin. Disaggregation into much smaller units of about 40,000 molecular weight occurs at high and low pH values or on treatment with various agents.

Earlier electron microscope studies by Hall ⁷⁵ revealed particles about 145 Å wide and 80 Å high which were obtained from dilute L-glutamate dehydrogenase (GDH) solution. Valentine ⁷⁶ observed spherical particles of 125 Å diameter, 90 Å and 75 Å in negatively stained GDH preparations. Although the objects varied in size, depending on the concentration of the protein, these electron micrographs were consistent with the concept of the undissociated molecule as a tetramer. Subsequent observations by Horne and Grevel ⁷⁷ indicate that the molecules tend to dissociate into typical tetrahedral subunits. However, attempts to observe the undissociated molecules have been hampered by technical difficulties.

Our recent investigations of bovine L-glutamate dehydrogenase , using micro-droplet cross-spraying techniques and micro-beam illumination for high resolution electron microscopy, have made it possible to observe the ordered aggregation and fine structure of the relatively well preserved molecules in the thin microcrystals. As shown in Figures 6 A,B,C,D, individual spherical or polyhedral particles of diameter 130 to 150 $\stackrel{\Omega}{ ext{A}}$ can be frequently observed, either free (Figure 6 C.D) or closely packed in ordered aggregates (Figure 6B) of the thin crystalline strands. The individual polyhedral particles of diameter 130 to 150 A may correspond to the undissociated individual molecules, while the numerous subunits of triangular shape could correspond partly to the GDH tetrahedral subunits described by Horne and Grevel. 77 However, on the basis of available data, the organization of the undissociated molecules appears to be more complex than the suggested simple aggregate of four subunits. Further studies are being pursued in attempts to elucidate the detailed subunit structure of these enzyme molecules. Investigation of the postulated changes in enzyme activity which may be produced by alterations of the subunit conformation of these enzyme molecules can also be of significance in relation to fundamental controlling mechanism in cells. Moreover, this enzyme is also of interest as a model system to illustrate the different stages of molecular and supramolecular organization in the observed paracrystalline arrays, which are relevant to the general problem of the organization of enzyme complexes in membranes.

C. Fraction-I Protein

We wish to refer to another example of the characterization of the substructure of a protein enzyme, which can be achieved by combined electron microscopic and biochemical studies. Fraction-I protein, which is a major soluble protein fraction located predominantly in the chloroplast and which comprises at least 50% of the soluble proteins, has marked carboxy-dismutase activity. Based on recent studies carried out in collaboration with Haselkorn, Kieras, and Van Bruggen, 79 Fraction-I protein of Chinese cabbage leaves is seen to consist of uniform cubical particles (Figure 7A) with an edge of about 120 Å.

In high resolution electron micrographs of negatively stained preparations (Figure 7B) the cubical particles exhibit a characteristic subunit structure which is consistent with a model having 24 subunits, in agreement with the present physical and chemical data. In the proposed model, the subunits may undergo considerable conformational changes, and do not appear to be related by principle of quasi-equivalence which governs the structural organization of many of the regular viruses. The fact that the substructure of a protein enzyme of only 120 Å can now be directly observed in electron micrographs, is of particular relevance to investigation of the enzyme complexes of similar size, which are presumably associated with nerve membranes and derivatives.

Membrane Ultrastructure in Nerve Cells - H. Fernández-Morán Subunit Patterns of Neuronal Membranes

A. Structural Subunit Patterns of Synaptic Membranes Investigation of the molecular organization of the synapse is of particular significance in view of the specialized molecular switching componentry assumed to mediate information storage, transfer and retrieval at these junctional regions.

Unfortunately, neuronal membranes have not yet yielded to correlate biochemical and electron microscopic studies of the type successfully applied to mitochondrial and chloroplast membranes. Difficulties in reliably isolating and preparing these labile structures without introducing serious artifacts impose serious limitations.

Typical subunit structural patterns in certain kinds of synaptic membranes have now been revealed by Robertson. 47

His comprehensive electron microscopic studies demonstrate a subunit pattern in the synaptic membranes of club-endings of Mauthner cells of the goldfish. Each of the alternating synaptic discs shows an internal beading with a repeat period of 90 Å to 95 Å in vertical sections. Transverse densities or lines spaced at 90 Å are seen in oblique sections, while frontal views of the membranes reveal a honeycomb pattern of lines and dots, approximately a hexagonal array. This network features dense granules in the center of each subunit facet and fine rodlet borders less than 20 Å in diameter. The resemblance of these patterns to the structure of Buckminster Fuller's geodesic domes suggests that this

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran type of efficient design may generally be embodied in the organization of cell membranes and regular virus shells. 47,73

Similar synaptic subunit structures have been seen in the Mueller cell of goldfish medulla. 47,80 Confirming and extending our earlier results,6,21 Robertson has demonstrated the existence of a regular subunit structure in the lamellae of retinal rod outer segments and in erythrocyte membranes. 80 This subunit pattern, which can be more readily seen when two membranes are closely apposed, is therefore, regarded as a structural pattern of general significance. 6,15,80

Robertson assumes as a working hypothesis that this pattern represents a structural protein on the outer surface of the membrane. Ocertain transverse densities which give the impression of a transverse globular substructure of the membranes are regarded with reservations since they may prove to be artifacts. However, only further work with improved preparation techniques and appropriate x-ray diffraction analysis can establish or preclude the existence of such a regular subunit organization in native undegraded neuronal membranes.

B. Organization of Synaptic Membrane Complexes

Several interesting ultrastructural features have been revealed in synaptic membrane complexes of different types. 80,81 Gray 81 describes in mammalian spinal cord ventral horn cells synaptic membrane complexes which exhibit 500 % dense particles

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran

in regular hexagonal array with a 1,000 A period. Certain preparation techniques reveal presynaptic fibrils and delicate hook-like fibrillar extensions from the postsynaptic cytoplasm, making contact in the synaptic cleft. In pyramidal cell synapses a patched differentiation of the synaptolemma is observed with two types of interlemmal elements which repeat at periods of about 235 A.80 An unusual subsynaptic formation has been described by Taxi80,81 in autonomic ganglia of certain amphibia. This complex is characterized by subsynaptic dense bands and granular regions forming a triangle with the subsynaptic membrane as its base. Identification and isolation of the specific macromolecular componentry subserving the complex processes at these junctional regions appears to be one of the most worthwhile goals requiring the full repertory of ultrastructural, chemical, and physiological investigations.

C. Fine Structure of Isolated Neuronal Membranes

The finest nerve fibers and neuronal extensions are fortunately of truly paucimolecular dimensions and, therefore, ideally suited for direct examination by electron microscopy. Confirming and extending our earlier work, 11,30,82 certain submicroscopic nerve fibers were found in frog spinal cord usually attached to the thicker nerve fibers. These fibers are thin enough to display their entire structure in a single electron micrograph. As shown in Fig. 8 A, these fibers appear as thin ribbons with diameters ranging between 0.1μ and 1μ and reaching lengths up to 50μ to 100μ .

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran

Essentially, each fiber as displayed in a whole-mount preparation, consists of a thin sheath formed by a single tubular membrane which contains long filaments 100 to 200 Å in diameter. In our original shadowed and osmium-fixed preparations no definite indications of fine, regular structure were observed in these membranes. However, with improved techniques new ultrastructural detail (Fig. 8 B) was revealed in these well-preserved, ultra-thin nerve membrane specimens. They exhibit a characteristic surface structure featuring in certain regions hexagonal arrays of closely packed polygonal elements 50 to 90 Å in diameter. Various type of dense granular components are found attached to the membranes, particularly in junctional regions as shown in Figs. 8 B and 9 A.B.

These techniques can undoubtedly be considerably refined, and it should eventually be possible to examine these ultrathin submicroscopic nerve fibers in vacuum-tight microchambers under conditions approaching the native hydrated state.

The Role of Water in Membranes

Water is a major constituent of nerve membranes, representing at least 35% to 40%, even in condensed multilayered, fluid-crystalline systems like myelin. The water layers at the aqueous interfaces of the fundamental repeating unit in myelin are about 12 $^{\rm A}$ to 15 $^{\rm A}$ thick. Their thickness appears to be largely determined by the electrical charge

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran density at the interfaces, and by the ionic strength. 2,3

Depending on the different environments we may envisage various types of water structures including water within the membrane structure itself, water in direct contact with the ordered macromolecular membrane surfaces, and water layers between closely paired systems of unit membranes. 91 Interconnected water channels within the myelin layers and related lamellar systems play an important role as possible pathways of diffusion of ions and certain molecules between the axon and extracellular fluids. 3

On the basis of results derived from a broad interdisciplinary approach \$^{91,98}\$ we must begin to consider water as a structured matrix, "the matrix of life" as was first pointed out by Szent-Györgyi. Substantial experimental evidence is now available in support of a mixture model to account for the unique properties of water. Thus, from neutron scattering data and other studies of water we are now able to envisage a sample of liquid water as being made up of "flickering clusters" of about 50 to 100 molecules with an average lifetime of the order of 10⁻¹¹ seconds, which is long enough for a structure to have a distinguishable existence.

The concept of organized water as an integral structural component of biological membranes must be given serious consideration. 2,6,88 We must assume that ordered water will specifically become integrated in the highly organized three-

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran

dimensional macromolecular structures of living systems. The abundance of postulated bulky species of water offers numerous possibilities, ranging from the crystalline hydrate structures to other frameworks of approximately tetrahedral fourfold coordination.

Based on our earlier correlated studies of water using nondestructive methods, such as nuclear magnetic resonance^{2,4,34} and x-ray diffraction techniques,^{4,14} the concept was developed of a three-dimensional hydrated lipoprotein system which provides a general structural framework for specialized macromolecular repeating units periodically arranged in the membrane layers.^{4,5,6,15} Although the main functional differences between membranes would depend upon the types of transducing units present and the complementary stereospecific configuration of the underlying substrate, this hydrated lipoprotein matrix could introduce a common unifying feature. The water, particularly in close association with the ordered lipoprotein membrane systems, must likewise be highly ordered, resembling "ice-like" hydration sheaths or crystalline hydrate lattices.

The open structure of water favors the formation of these crystalline hydrates ⁹⁴ which are clathrate compounds with a hydrogen-bonded, ice-like framework defining cavities able to enclose molecules of noble gases, liquids, ions, protein side chains, etc., of appropriate size (4 to 6.9 Å). The hydrogen-bonded lattice of these microcrystalline hydrates, which are about 12 Å to 24 Å, is stabilized by van der Waals

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran

interaction, in contrast to ordinary ice. The hydrate crystals may therefore remain stable even in the range of normal body temperatures under specialized conditions. As suggested earlier, 5,6,87,88 these polyhedral hydrate structures with numerous built-in cavities could effectively enclose ions, or form closely fitting "molecular replicas" around protein side chains, thus adapting to the specific macromolecular configurations by three-dimensional interpenetration.

Crystalline hydrate structures of this general type, or equivalent organized water structures permeating the ordered lipoprotein-subunit macromolecular assembly system would provide an interconnected H-bonded substrate for fast protonic charge-transport mechanisms.

Localized reversible phase changes in such ordered water structures could also provide the basis for a variety of conformational changes in protein and lipoprotein layers, including reversible transitions from coherent bimolecular lipid layers to the subunit micellar arrays. Reversible phase transitions of this kind, spreading through the membrane matrix, may contribute significantly to the propagation of local perturbations. Likewise, selective permeability may be envisaged in terms of molecular "pores" lined with ordered water. In general, this concept of ordered water lattices forming an integral structural component of membranes provides a powerful conceptual basis for understanding many fundamental mechanisms of membrane function, 91,98 as pointed out by Hechter.

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Morán

The precise molecular nature of the ordered water in membranes is, however, still unknown. Nevertheless, nuclear magnetic resonance techniques have been successully applied in studies on the hydration structure of fibrous macromolecules. 99

Despite the uncertainties involved, there is recent evidence supporting the structural role of water in membrane systems. For example, Person and Zipper 91 have shown that treatment of mitochondrial suspensions with a dry zeolite, which clathrates the water, produces the same type of membrane disruption as is usually achieved with detergents. This novel form of membrane disruption can, therefore, be attributed to the avidity of the zeolite preparation for water, and its capacity to selectively remove certain divalent cations.

Extending our earlier work, ⁸⁸,92,93 we have tried to obtain direct experimental verification of organized water structures of the crystalline hydrate type through controlled local formation of microcrystalline, noble gas hydrates in selected lipid, lipoprotein complexes, and cell membranes. ⁹⁸ As shown in Figure 10 B characteristic electron-dense microcrystalline hydrates, about 10 Å to 20 Å in diameter, are localized mainly in the hydrophilic regions of lipid micelles following application of argon and xenon under controlled high pressure and temperature in special specimen chambers. The noble gas atoms hereby occupy polyhedral cavities in a hydrogen-bonded framework of water molecules, contributing to stabilize and make electron optically visible any pre-existing organized water structures of the hydrate type in lipid or lipoprotein model systems.

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran

These techniques for direct visualization of noble gas hydrates could eventually provide experimental data bearing on the existence of the hydrate microcrystals reversibly forming in cell membranes. This approach is relevant to the molecular theory of general anesthesia postulated by Pauling. According to this theory the formation of clathrate-like structures by anesthetic agents, including noble gases, in the aqueous portion of nervous tissues could cause anesthesia by modifying the electrical activity at junctional regions through entrapment of ions and electrically charged side chains of protein molecules. The formation of these hydrates would increase the structured water in the membrane systems, thus presumably making the water at synapses less available for facilitating conduction.

One might even go beyond this theory of Pauling, and in common with the earlier hypotheses of anesthesia advanced by Claude Bernard, 96 postulate reversible phase changes or "coagulations" in the synaptic membranes. It might not be unreasonable to assume that these reversible transformations in nerve membranes are not only associated with anesthesia, but perhaps also with certain rhythmic changes such as sleep.

Reversible modifications of the molecular organization involving primarily the hydrated lipoprotein framework of nerve membranes must, therefore, ultimately take into account the structure and cooperative ordering of water.

MEMBRANE ULTRASTRUCTURE IN NERVE CELLS - H. Fernández-Morán The Biosynthesis of Membranes

The general problem of membrane biosynthesis is of fundamental importance for all biological studies concerned with embryogenesis, morphogenesis, and differentiation.

The available evidence indicates that characteristic mechanisms are involved in the biosynthesis of the protein and lipid membrane components, and also suggest the concept that membranes may furnish the template for their "self-duplication". 109,110

Investigation of nerve membrane biosynthesis may prove to be of particular value in elucidating the synthesis and assembly of the repeating subunits and their organization within the plane of the membrane layers. The nature of the association of the protein synthetic machinery and of nucleic acids with cell membranes is therefore being studied by electron microscopy and biochemical techniques to serve as a basis for a better understanding of membrane biosynthesis.

A. <u>Mitochondrial DNA, RNA, and Protein Biosynthesis System</u> in Membrane Replication.

The discovery of characteristic nucleic acids in mitochon-dria 101,102,103 and demonstration of a protein biosynthesis system have recently confirmed and extended earlier work

(102) to provide significant support for the semi-independence or autonomy of these organelles. DNA of mitochondria from various mammalian cell types 103,104 differ from nuclear DNA in guanine plus cytosine content, and in their renaturation properties. In contrast to nuclear DNA which is linear, the molecules of mitochondrial DNA exist in circular form, their mean contour lengths measuring 5.24µ-5.45µ. Two to six ring molecules of molecular weight 10 x 106 may be present in a single mitochondrien. Attachment of membrane pieces to DNA, and the observed association of mitochondrial DNA with cristae may possibly represent sites where replication is initiated. Recent evidence also indicates that mitochondria contain mechanisms for incorporation of nucleotides into this DNA.

Accumulating evidence indicates that in addition to DNA, RNA, ribosome-like particles 102, and RNA polymerase-like systems 66,102 have been found in mitochondria from different sources.

This new information furnishes the basis for a working hypothesis implying that mitochondria possess their own genetic information and the necessary mechanism to express this information in an autonomous synthesis of proteins. 66 A 5.5µ long DNA molecule is able to code the information for the synthesis of several hundred proteins. As pointed out by Andre, part of this information may function as structural

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran genes, controlling the duplication of mitochondria and the synthesis of the biochemical equipment. Another part could function as regulatory genes, controlling differentiation and response to the external medium. Further studies will help to establish to what extent these genes are hypothetical, and to what extent they are subordinate to information coming from the nucleus, or may be reacting independently to the external medium by mutation and adaptation .66

These hypothetical considerations may prove to be of operational significance in view of the characteristic association of mitochondria with synaptic endings. 80,81 In monkey spinal cord motor horn cells Bodian has described very large endings containing unusually densely packed mitochondria. The subsynaptic cytoplasm in many of these large endings is characterized by an accumulation of Nissl substance which consists of regular layers of endoplasmic reticulum cisternae with large numbers of ribosomes in between. It has been speculated that these Nissl bodies may be associated with protein synthesis in relation to specific endings. However, in view of the dense accumulation of mitochondria possessing their own genetic information and the mechanisms to express it in an autonomous synthesis of protein, the interesting possibility of a more direct participation of mitochondria in regulatory aspects of these biosynthetic processes should be taken into consideration. Subunits of protein molecules would be synthesized locally and either be attached as specific

Membrane Ultrastructure in Nerve Cells - H. Fernández-Moran macromolecular assemblies corresponding to coding and information storage in molecular switching componentry at the junctional regions, or they might pass through the ER membrane and the cysternal cavity into the extracellular space. Similar considerations might also apply to the intimate interrelationship of mitochondria and associated granular components with photoreceptors of the insect retinula as shown in Fig. 11.

B. Role of RNA Polymerase in transfer of Genetic Information. The enzyme RNA polymerase plays a key role in the transfer of genetic information through its participation in the differential RNA transcription upon DNA templates. 106,107 Confirming the results of earlier workers we have studied the physical properties of DNA-dependent RNA polymerase known to be relatively free of nucleic acid and capable of asymmetric transcription.

As shown in Fig. 12 b, the RNA polymerase molecules from $\underline{\mathbf{E}}$. $\underline{\operatorname{coli}}$, prepared by the Chamberlin and Berg procedure, and having a sedimentation coefficient of 25 s, appear to consist of six subunits arranged in a hexagon with a cross section of 120 to 130 $^{\mathrm{A}}$ in negatively stained preparations. If the large hexagonal structures are the RNA polymerase molecules, as proposed by Fuchs, Zillig et al, the basis structure is either a hexagonal disc or two of these stacked together. A few smaller structures are also observed which might be degradation or dissociation products, some of which closely resemble the subunits in the hexagon. As shown by J. P. Richardson, 107 the ionic conditions strongly affect the form of the RNA polymerase.

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran

The low salt conditions favorable for observing the 21 s form are the optimal conditions for enzymatic activity, while the high salt conditions which produce the 13 s form completely inhibit RNA polymerase activity.

In general, these observations emphasize the critical effect of ionic conditions on all important systems participating in the process of biosynthesis. Bearing this in mind, one is tempted to speculate on the possible effect of ionic currents which might have been modulated by bioelectric potentials at key junctional membrane regions, thus presumably affecting the processes which regulate information transfer and storage in nerve membranes and their derivatives.

In order to obtain more information on the manner in which RNA polymerase initiates transcription of RNA upon DNA templates, we are carrying out correlated electron microscopic studies on the binding of RNA polymerase to different types of DNA. Following the work of H. S. Slayter and C. E. Hall, 106 and J. P. Richardson 107 it is assumed that certain sites must exist along the DNA molecules which specifically bind RNA polymerase, allowing transcription to begin at well-defined regions. Preparations of RNA polymerase were mixed in with different types of DNA, including \emptyset X 174 DNA, bacterial and salmon DNA, in varying concentrations under carefully controlled conditions.

As shown in a representative example (Fig. 12 a) the complexed

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran enzyme generally appears to center itself upon the DNA axis, which is in agreement with the results reported by Slayter and Hall. Distribution of the RNA polymerase molecules along the DNA fibers varied according to the different preparative conditions. However, the disposition of enzyme particles is similar to the distribution reported by Slayter and Hall, with average separations ranging from about 1,200 A to about 2,000 Å. These preliminary results are in agreement with values reported by other workers. However, further studies with improved preparation techniques are required before a reliable evaluation can be carried out. Particular interest attaches to correlated studies on the binding of RNA polymerase to circular forms of DNA (e.g. from Ø X 174) which are being carried out in collaboration with Dr. Samuel Weiss and his associates.

C. Morphopoietic Genes and Membrane Biosynthesis.

As indicated earlier, the physical principles underlying the designs of regular virus shells 73 may be applicable to cell membranes assuming that the constituent protein or lipoprotein subunits are organized in a lattice structure. When dealing with the simplest virus shells we can envisage that the subunits have complete shape-specificity, and thus contain all the information needed for self-assembly of identical subunits required to build up the prescribed shape.

More complex shells require additional information which is stored in morphopoietic genes. 122 In T4 bacteriophage the

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran functions of three such genes are known, which appear to require a structural scaffolding by means of which this information is expressed.

There is reason to believe that we will have to cope with a major problem in this area when trying to understand the organization and functioning of cell membranes and their derivatives, which represent the next higher order of molecular structure in cellular organization. Knowledge of the genetic control of shape in protein structures will probably not be enough to explain the origin of the complex membrane-bound shapes of the highly specialized cells in the nervous system. However, in view of the complex relationships involved in membrane morphogenesis and differentiation, high resolution electron microscopy may be expected to play an important role in direct visualization of the sequential molecular patterns involved in nerve membrane biosynthesis.

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran

Cooperative Features of Nerve Membranes.

Biological membranes are the largest and most complex atomic aggregates known, spanning the widest range of dimensions as basic integrating components of all living systems. Elucidation of the molecular organization of cell membranes represents, therefore, not only one of the major fundamental problems of biomedical research, but it has also emerged as the next major barrier, cutting across molecular biology and neurobiology, which must be overcome to clear the way for further progress.

Membranes exhibit a unique degree of structural asymmetry, comprising a coherent lipoprotein film that is usually only a few molecules thick, but still capable of extraordinary lateral extension.

Nerve membranes exhibit these characteristics to a unique degree, as major components of the nervous system, which is generally regarded as the most sophisticated and complex biological system known. Thus, the total length of the peripheral network of the human nervous system is estimated at about 600,000 miles. All fibers of the human body's nerve fabric, which is essentially lined with multiply folded membranes, would extend three times the distance from the earth to the moon. Having unravelled their exquisitely delicate, yet highly ordered texture, we cannot help wondering about the nature of these "paucimolecular" "living" films, singularly sensitive to "paucienergetic

Membrane Ultrastructure in Nerve Cells - H. Fernández-Morán stimuli", 108 which are so compactly infolded into the submicroscopic recesses of the relatively limited space occupied by the individual nervous system, that we could encompass a whole planet with the combined array of extended membranes.

In many respects, the entire complex membrane repository comprising the nervous system may be regarded as representing an essentially interconnected, continuous and "closed membrane system" where all junctional domains of the separate cell membrane entities establish functional contact and are interrelated to form an integrated, coherent whole of extraordinary dimensions. In this sense, if a diamond crystal may be considered as one giant "macromolecule" so it is justified to conceive of these CNS membrane aggregates as one giant "megamembrane" or "megamolecule". Such a coherent system would be uniquely suited to effect transfer, storage and retrieval of information at the various hierarchies of organization.

Although this model is strongly idealized, it would, nevertheless, embody the basic features to serve as a suitable substrate for highly cooperative phenomena embracing the entire complex molecular aggregates, which are not unlike those involved with manifestation of quantum phenomena at the macroscopic scale, like for example, in superconductivity. Although admittedly speculative and only intended to delineate salient features of a model, this approach may nevertheless prove to be a convenient working hypothesis which could provide the operational key to a deeper under-

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran standing of the molecular and supramolecular organization features of the CNS.

Macromolecular Repeating Units and Domains in Membranes.

There is another feature of nerve membranes, which although derived from the results of ultrastructure research, nevertheless, attempts to correlate them with the overall behavior of the system. Thus, membranes may also be regarded as arrays of organized domains or subunits, including macromolecular assemblies. Katchalsky, et al 23 conceive of such cooperative systems as exhibiting "memory-hysteresis behavior". It is pointed out that there exists also the possibility of a physical record in biopolymers based on conformational changes in single macromolecules or in cellular macromolecular structures such as membranes. Typical examples of physical memory devices are the magnetic tape of tape recorders. Katchalsky conceives of the possibility that conformational transitions in macromolecules might also exhibit hysteresis phenomena. Also, the existence of various distributions of functional side-groups along the macromolecules makes possible the existence of different domains with a wide range of transition points. 123 Metastable macromolecular states of high permanence are linked with two-dimensional hysteresis phenomena based on domains in biological films.

According to Katchalsky et al, 123 if we assume that the smallest number of nucleotides is about 8-10 in a domain,

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran

the energy involved in a phase transition of a nucleotide is higher than 2 RT, which is a plausible figure in case memory recording in domain structures is considered. This figure is sufficiently high to make a signal changing the domain structure distinguishable from the thermal noise, but yet small enough to be of use in the low-energy transformations of living organisms. In this connection, it is pertinent to point out that membranes as two-dimensional ordered arrays of macromolecular assemblies, acting like information storage devices, would ideally fulfill the dual criteria of "low energy switching", and optimum size of domains as metastable forms of high permanence.

Cooperativity of Membranes and their Subunit Structure.

Review of the data on membrane organization as revealed by correlated ultrastructural and biochemical studies discloses certain general features. Coherent paucimolecular layers of indefinite lateral extension appear to consist of a periodic, hydrated lipoprotein substrate which is integrated with specific macromolecular repeating subunits organized in asymmetric, "paracrystalline", arrays within the plane of the layers. There is a fundamental underlying concept that membranes are made up by the stereospecific association of repeating macromolecular subunits, with specific conformations dependent on their association with the membrane substrate.

Recently, J. P. Changeux and his colleagues 132 have extended their theory of allosteric transitions in enzyme-substrate

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran reactions 125 to membranes composed of identical units, applying some of the principles and approaches proposed for regulatory enzymes. They discuss the cooperative properties of membranes on the basis of their highly ordered structure.

Based primarily on data derived from correlated ultrastructural and biochemical studies of membranes, carried out during the past years by Fernandez-Morán, Green, Perdue, et al, 6,15,17 they consider the following significant aspects of membrane organization: (a) membranes are made up by the association of repeating macromolecular lipoprotein units; (b) the conformation of these units (protomers) differs when they are organized into a membrane structure or dispersed in solution; (c) many biological or artificial lipoprotein membranes respond in vivo, as well as in vitro, to the binding of specific ligands by some modification of their properties which reflects rearrangement of the membrane organization and presumably of the repeating unit's conformation.

From this they deduce a theoretical model of a biological membrane as an ordered collection of repeating globular lipoprotein units, or "protomers" organized into a two-dimensional, crystalline lattice. The protomer constitutes the "primitive cell" of the lattice and does not necessarily possess in itself any particular symmetry properties. Several conformational states are reversibly accessible to the protomer. The conformation of the protomer depends upon

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran

its association with neighboring protomers, and is thus submitted to a lattice constraint similar to the quaternary constraint involved in the organization of the quaternary structure of oligomeric proteins. This interesting theory has been primarily developed to account for the cooperative phenomena accompanying the binding of ligands to a membrane. This model predicts the two classes of responses exhibited by biological membranes: a "graded" response or an "all-or-none" response. Basic to their theory is the assumption that "conformational changes of the protomer not only preexist the ligand binding but are not fundamentally different whether the ligand is bound or not." These authors emphasize that a number of important biological phenomena seems to be related to the highly cooperative structure of cell membranes.

<u>Multiplex Systems and Possible Ultrastructure Correlates</u> <u>in Nerve Membranes</u>.

There appears to be a deep analogy between von Neumann's multiplexing concept 124 (which essentially introduces redundancy so that the reliability of the whole system is greater than the reliability of its parts) and the hierarchically organized redundancy of structural substrates in the CNS at the macromolecular and molecular levels now revealed by electron microscopy. The microcomponentry required by the multiplexing technique may partly correspond to the immense number (in the order of ca. 10¹⁶ to 10¹⁹) of specific macromolecular assemblies and related subunits periodically arranged on the nerve membranes and derivatives

Membrane Ultrastructure in Nerve Cells - H. Fernández-Morán in close association with axon filaments, ribosomes, and other key components.

Major Problems in the Study of Nerve Membrane Ultrastructure. With the improved methodology now available it is tempting to systematically extend the fine structure studies to all regions of the nervous system in vertebrate and invertebrate specimens. 126 However, one should bear in mind that this purely descriptive approach could easily consume the best efforts of entire generations of investigators. Thus, even if perfect techniques were currently available, a complete morphological analysis of three-dimensional relationships based on electron microscopy of serial sections would involve such a prohibitively large number of images to map only a few cubic centimeters of nerve tissues that total man hours would figure in the order of centuries.

Instead, it appears more profitable to concentrate on certain key problems of nerve ultrastructure and function:

(a) Correlated biochemical, electron microscopic and biophysical characterization of the major protein, lipoprotein, glycoprotein components, and the macromolecular and multienzyme assemblies associated with nerve membranes in a

Membrane Ultrastructure in Nerve Cells - H. Fernández-Morán variety of specimens, including human. Attention would center on well-characterized enzymatic and protein components with specific physiological properties. 127

- (b) Systematic studies of the organization of subunit patterns within the plane of membrane layers, preferably examined in the native hydrated state under conditions of reduced radiation damage and of minimum perturbation, by using improved instrumentation and preparation techniques for high resolution, low-temperature electron microscopy correlated with polarized light and x-ray diffraction studies. 2,6,33,92,93,98, 113,114,116
- (c) Correlation of ultrastructural studies with a comprehensive "macromolecular and molecular anatomy program" aimed at achieving reliable dissection and quantitative separation of the entire spectrum of discrete molecular species using Anderson's zonal centrifuge techniques 133 and related methods.
- (d) Correlative studies of specific, dynamic changes on selected nerve specimens in vivo 129 and in vitro 130 (tissue and organ cultures) using light and electron microscopy.

It would be of special interest to carry out a correlated ultrastructural and biochemical analysis of the molecular organization of selected key sites of the central nervous system by performing a series of "submicroscopic or macromolecular biopsies". Such a submicroscopic biopsy, for example of key synaptic sites or cerebral nuclei, can be

carried out "in vivo" using precision stereotactic techniques under conditions of minimum perturbation without inflicting more than "subliminal" damage, which is perhaps hardly greater than the natural rate of decay of nerve cells taking place in the adult brain. These submicroscopic CNS biopsies, carried out with specially developed ultramicrocapillaries, can be combined with subsequent tissue culture explants and related in vitro techniques. Hyden has already carried out significant pioneering work in microbiopsies of living tissues. Further development of these techniques holds great potential for electron microscopy.

Ultrastructural studies could also be combined with appropriate non-destructive techniques, including nuclear magnetic resonance, ³⁴ electron spin resonance, and methods for investigation of fast reactions. ¹⁰⁰ Neutron diffraction and scattering studies ⁹⁷ suitably modified for examination of nervous tissues should also yield valuable information.

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran

Suggested Methodological Approaches in Correlated Studies of Nerve Membranes.

Recent improvements in instrumentation and preparation techniques for electron microscopy appear to be especially suited for correlative investigations of nerve membrane ultrastructure at the molecular and pauciatomic levels 116: (a) High resolution phase contrast electron microscopy with improved point cathode sources, low temperature stages and specially stabilized power supplies, as well as new types of phase contrast apertures has enabled us to visualize directly for the first time (Fig. 13b) the hexagonal cells of single crystal graphite containing only 10 carbon atoms. Demonstration of this capability of pointto-point resolution in the range of interatomic distance now makes it appear feasible to record detail of molecular and pauciatomic dimensions in biological specimens. Extension of this work and the enhanced capability of interpreting phase contrast images should eventually enable us to achieve direct readout of molecular and submolecular structures, particularly in nerve membranes. Parallel advances in lowtemperature electron microscopy at liquid helium temperatures using specially developed superconducting lenses, 113,114,116c intensifiers and improved recording techniques should image eventually permit us to see directly the three-dimensional structure of key regions of synaptic junctions and of other nerve membrane areas.

Membrane Ultrastructure in Nerve Cells - H. Fernández-Morán

- (b) By using vacuum-tight microchambers 92-98 with ultrathin windows and related techniques, it is possible to observe membranes under conditions approaching their native hydrated state. Examination of living hydrated specimens can now be carried out thanks to recent advances in high voltage electron microscopy. 115 Larger microchambers for examination of intact, viable biological specimens have been successfully applied owing to the far greater penetration now possible with advanced high voltage electron microscopes. Moreover, by using pulsed T-F emission or stroboscopic microbeam illumination, one can, in principle, obtain a unique combination of high spatial and high temporal resolution for the study of fast reactions in nerve membranes.
- (c) Scanning electron microscopy can likewise be profitably used in the study of nerve membrane specimens examined under favorable conditions. As shown by the recent remarkable work of R. F. Pease et al, 131 electron micrographs of living specimens of various developmental stages of the insect Tribolium confusum have been obtained with a scanning electron microscope at magnifications of about 1,000 X. The requisite conditions of moderate vacuum and the relatively minor effects of irradiation by the imaging electron beam, which have been shown to be compatible with survival of certain specimens, may be applied in suitably designed experimental chambers to study selected nerve membranes and nerve cell specimens. Scanning electron

Membrane Ultrastructure in Nerve Cells - H. Fernández-Morán microscopy can also be extended to the application of special bioelectric and biomagnetic probes for the study of rapid and ultrarapid phenomena in nerve membrane systems.

- (d) We can also foresee the unique potential of electron microscopy "applied in reverse" for achieving electron optical demagnification of letters, diagrams, printed circuits, etc., in such a way that in the optimal case, the contents of a several million volume library can be condensed on an area of a single page size or its equivalent. 117,118
- (e) The same electron optical techniques which revealed the exquisite molecular componentry of nerve cells may some day serve to duplicate true homologous ultraminiaturized, "molecular computer and information storage systems." Although the packing density embodied in the human brain is of the order of 10⁹ components per cubic inch, impressive advances made during the past years in manufacturing integrated microelectronic systems have already resulted in practical packing densities of about 10⁵ elements per cubic inch, which is susceptible of further improvement. Likewise, functional printed circuits of micron or submicron dimensions which could fit on a red blood cell, for example, and transmit important functional data recorded <u>in vivo</u> at the cellular and subcellular levels are within the scope of present technology.

One can conceive of integrated ultramicroelectric circuits which begin to approach the dimensions of macromolecular assemblies, and could eventually be incorporated

Membrane Ultrastructure in Nerve Cells - H. Fernández-Morán into key junctional sites of living nerve membranes without undue perturbation. Such submicroscopic prosthetic sensors, with their envelopes composed of biosynthetically, produced macromolecular "protein coats" to form integral components of the nervous systems, could well subserve certain unique functions. For example, since they can be produced in large quantities and implanted in sufficient numbers throughout the central nervous system, they can provide a direct operational link at the crucial macromolecular level between the central nervous system and man-made information processing systems, such as computers of commensurate complexity.

There is increasing evidence that biological membrane systems may have properties in common with semiconductors. 2,119,120 It may therefore be of interest to reconsider as a working hypothesis suggestive of new experimental approaches our earlier speculations on certain specific crystalline properties such as piezoelectric effects, semiconductor properties or equivalent phenomena which may be associated with the fluid-crystalline nature of myelin and related lamellar systems. 2,30 In connection with correlated studies on basic mechanisms of electoencephalography, I had previously discussed the hypothetical possibility that myelin and other paracrystalline nerve membrane systems might function as generators or specific amplifiers of coherent electromagnetic radiation, mainly in the infrared or UHF range between 1 and about 10 microns wavelength. 134 hypothetical mechanisms involving far infrared quanta and

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran

coherent phenomena in the transfer of biological information on the macro- and supramolecular level are now being considered by J. Polonsky 135 in the scope of molecular exciton theory and quantum electronics.

Conclusions

Obviously, our current knowledge about nerve membrane ultrastructure and function is but a presage of many as yet unknown and unexpected properties of these basic constituents of the nervous system, awaiting further investigation.

Ultrastructure research should eventually enable us to learn how all of the prodigious activity of nerve cells and their appendages may be unfolding invisibly at the molecular and atomic levels in a strange, spatial-temporal framework, possessing an order that one may some day aspire to comprehend.

The old adage may thus prove to be invested with a deeper meaning when the simile is invoked that the neurons apparently have so little to show for all their incessant submicroscopic labors, because, like the mills of the gods, they grind exceedingly small--between membrane interfaces at the molecular level.

REFERENCES

- Fernandez-Morán, H., and Brown, R., 1958. The submicroscopic organization and function of nerve cells, Exptl. Cell Research, Suppl., Vol. 5, p. 1.
- Fernández-Morán, H., 1959. Fine structure of biological lamellar systems, in Biophysical Science -- A Study Program, (J. L. Oncley, et al, editors), John Wiley & Sons, Inc., New York, and Revs. Modern Phys., Vol. 31, pp. 319-330.
- 3. Schmitt, F. O., 1959. Molecular organization of the nerve fiber, in Biophysical Science -- A Study Program, (J. L. Oncley, et al, editors), John Wiley & Sons, Inc., New York, and Revs. Modern Phys., Vol. 31, pp. 455-465.
- 4. Fernandez-Morán, H., 1961. Lamellar systems in myelin and photoreceptors as revealed by high-resolution electron microscopy, in Macromolecular Complexes, (M. V. Edds, Jr., editor), New York, The Ronald Press Company, pp. 113-159.
- 5. Fernández-Morán, H., 1962. Molecular basis of specificity in membranes, in Macromolecular Specificty and Biological Memory, (F. O. Schmitt, editor), Cambridge, Mass., The M.I.T. Press, pp. 39-48.
- 6. Fernández-Morán, H., 1962. Circulation, Vol. 26, p. 1039.
- 7. Schmitt, F. O., 1965. The physical basis of life and learning, Science, Vol. 149, pp. 931-936.
- 8. Schmidt, W. J., 1937. Die Doppelbrechung von Karyoplasma, Zytoplasma, und Metaplasma, Berlin, Gebruder Borntraeger.

REFERENCES

- 9. Schmitt, F. O., Bear, R. S., and Clark, G. L., 1935 X-ray diffraction studies on nerve, Radiology, Vol. 25, p. 131.
- 10. Fernández-Morán, H., 1950.. Sheath and axon structures in the internode portion of vertebrate myelinated nerve fibers, an electron microscope study of rat and frog sciatic nerves, Exptl. Cell Research, Vol. 1, pp. 309-340.
- 11 Fernandez-Morán, H., 1952. The submicroscopic organization of vertebrate nerve fibers, an electron microscope study of myelinated and unmyelinated nerve fibers, Exptl. Cell Research, Vol. 3, pp. 282-350.
- 12 Davson, H., and Danielli, J. F., 1943. The Permeability of Natural Membranes, Cambridge University Press.
- 13. Robertson, J. D., 1959. The ultrastructure of cell membranes and their derivatives, Biochem Soc. Symposia (Cambridge, England), Vol. 16, p. 3.
- 14. Fernandez-Morán, H., 1962. Cell-membrane ultrastructure: low temperature electron microscopy and x-ray diffraction studies of lipoprotein components in lamellar systems, in Ultrastructure and Metabolism of the Nervous System, (S. R. Korey, editor), Baltimore, The Williams & Wilkins Company, A.R.N. M.D. Series, Vol. 40, pp. 235-267.
- 15 Fernández-Morán, H., Oda, T., Blair, P. V., and Green, D. E, 1964. A macromolecular repeating unit of mitochondrial structure and function, J. Cell Biol., Vol. 22, pp. 63-100.

MEMBRANE ULTRASTRUCTURE IN NERVE CELLS - H. Fernández-Morán REFERENCES

- 16. Smith, D. S., 1963. The structure of flight muscle sarcosomes in the blowfly <u>Calliphora</u> erythrocephala (Diptera), J. Cell Biol., Vol. 19, p. 115.
- 17. Green, D. E. and Perdue, J.F., 1966. Membranes as expressions of repeating units, PNAS, Vol. 55, pp. 1295-1302.
- 18. Lehninger, A. L., 1964. The Mitochondrion, New York, W. A. Benjamin, Inc.
- 19. Park, R. B. and Pon, N. G., 1963. Chemical composition and the substructure of lamellae isolated from Spinacea oleracea chloroplasts, J. Mol. Biol., Vol. 6, p. 105.
- 20. Park, R. B. and Biggins, J., 1964. Quantasome: size and composition, Science, Vol. 144, pp. 1009-1011.
- 21. Fernandez-Moran, H., 1961. The fine structure of vertebrate and invertebrate photoreceptors as revealed by low temperature electron microscopy, in The Structure of the Eye, (G. K. Smelser, editor), New York, Academic Press, pp. 521-526.
- 22. Robertson, J. D., 1964. Unit membranes: a review with recent new studies of experimental alterations and a new subunit structure in synaptic membranes in Cellular Membranes in Development, (M. Locke, editor), New York, Academic Press, pp. 1-82.
- 23. Robertson, J. D., 1962. The unit membrane of cells and mechanisms of myelin formation, chapter VI in Ultra-structure and Metabolism of the Nervous System, Research Publications, Association for Research in Nervous and Mental Disease, (S. R. Korey, A. Pope and E. Robins, eds.)

 Baltimore, The Williams & Wilkins Company, Vol. 40, pp.94-158.
- 24. Branton, D., 1966. Fracture faces of frozen membranes, Nat'l. Acad. Sci., Vol. 55, pp. 1048-1056.

MEMBRANE ULTRASTRUCTURE IN NERVE CELLS - H. Fernández-Moran

REFERENCES

- 25. Overton, J., Eichholz, A., Crane, R. K., 1965. Studies on the organization of the brush border in intestinal epithelial cells, J. Cell Biol., Vol. 26, p. 693.
- 26. Benedetti, E. and Emmelot, P., 1965. Electron microscopic observations on negatively stained plasma membranes isolated from rat liver, J. Cell Biol., Vol. 26, pp. 299-305.
- 27. Cunningham, W., Stiles, J., and Crane, F., 1965. Exptl. Cell Res., Vol. 40, p. 171.
- 28. Schmitt, F. O., 1950. The colloidal organization of the nerve fiber, in Genetic Neurology (P. Weiss, ed.), Chicago, Illinois, University of Chicago Press, pp. 40-52.
- 29. Schmitt, F. O., 1958. Axon-satellite cell relationships in peripheral nerve fibers, Exptl. Cell Res., Suppl. Vol. 5, pp. 33-57.
- 30. Fernández-Morán, H., 1954. The Submicroscopic structure of nerve fibers, Progr. in Biophys. and Biophys. Chem., Vol. 4, pp. 112-147.
- 31. Geren, B. B., and Schmitt, F. O., 1955. Electron microscope studies of Schwann cell and its constituents with particular reference to their relation to the axon. In Symposium on Fine Structure of Cells (Congress of Cell Biology, 8th, Leyden, 1954), New York, Interscience Publishers, p. 251.
- 32. Finean, J. P., 1958. X-ray diffraction studies of the myelin sheath in peripheral and central nerve fibres, Exptl Cell Res., Suppl., Vol. 5, pp. 18-32.
- 33. Fernández-Morán, H., Finean, J. P., 1957. Electron microscope and low-angle x-ray diffraction studies of the nerve myelin sheath, J. Biophys. Biochem. Cytol., Vol. 3, pp. 725-748.
- 34. Fernández-Morán, H., 1957. Electron microscopy of Nervous Tissue, in Metabolism of the Nervous System, II International Neurochemical Symposium, Aarhus (July 1956), London, Pergamon Press, pp. 1-34.

MEMBRANE ULTRASTRUCTURE IN NERVE CELLS - H. Fernández-Morán

REFERENCES

- 35. O'Brien, J. S., 1965. Stability of the myelin membrane, Science, Vol. 147, pp. 1099-1107.
- 36. Folch-Pi, J., 1955. In <u>Biochemistry of the Developing Nervous</u>
 System (H. Waelsch, Ed.), New York, Academic Press, p. 121.
- 37. Vandenheuvel, F. A., 1963. In J. Am. Oil Chemists' Soc., Vol. 40, p. 455.
- 38. Waelsch, H., Sperry, W., and Stoyanoff, V., 1940. J. Biol. Chem., Vol. 135, p. 291.
- 39. Davison, A. N. and Dobbing, J., 1960. Biochem. J., Vol. 75, p. 565.
- 40. Singer, M. and Salpeter, M., 1966. Transport of tritiumlabelled L-histidine through the Schwann and myelin sheaths into the axon of peripheral nerves, Nature, Vol. 210, pp. 1225-1227.
- 41. Schmitt, F. O., and Davison, P. F., 1965. Role of protein in neural function, Neurosciences Research Program Bulletin, Vol. 3, pp. 55-76.
- 42. Huneeus-Cox, F., 1964. Electrophoretic and immunological studies of squid oxoplasmic proteins, Science, Vol. 143, pp. 1036-1037.
- 43. Huneeus-Cox, F. and Smith, B. H., 1965. The effects of oxidizing, reducing and sulfhydryl reagents on resting and action potentials of the internally perfused axon of Loligo pealeii, Biol. Bull., Vol. 129, p. 408.
- 44. Branton, D., 1966. Fracture Faces of Myelin, in press.
- 45. Moor, H., Personal communication.
- 46. Branton, D. and Park, R. B., Personal communication.
- 47. Robertson, J. D., 1963. The occurrence of a subunit pattern in the unit membrane of club endings in Mauthner cell synapses in goldfish brains, J. Cell Biol., Vol. 19, pp. 201-221.
- 48. Schmitt, F. O., 1963. The Macromolecular assembly—a hierarchical entity in cellular organization, Develop. Biol., Vol. 7, pp. 546-559.

MEMBRANE ULTRASTRUCTURE IN NERVE CELLS - H. Fernández-Morán REFERENCES

- 49. Sjøstrand, F. S., 1949. J. Cellular Comp. Physiol., Vol. 42, pp. 15-44.
- 50. Finean, J. P., Sjöstrand, F. S., and Steinman, F., 1953. Exptl. Cell Res., Vol. 5, p. 559.
- 51. Blasie, J. K., Dewey, M. M., Blaurock, and Worthington, 1965. Electron microscope and low-angle x-ray diffraction studies on outer segment membranes from the retina of the frog, J. Mol. Biol., Vol. 14, pp. 143-152.
- 52. McConnell, D. G., 1965. The isolation of retinal outer segment fragments, J. Cell. Biol., Vol. 27, pp. 459-473.
- 53. Fernández-Morán, H., 1956. Fine structure of the insect retinula as revealed by electron microscopy, Nature, Vol. 177, pp. 742-743.
- 54. Wolken, J. J., Copenos, J., and Turano, A., 1957. J. Biophys. Biochem. Cytol., Vol. 3, p. 429.
- 55. Fernández-Morán, H., 1958. Fine structure of the light receptors in the compound eyes of insects, Exptl. Cell Res., Suppl. 5, pp. 586-644.
- 56. Waterman, T. H. and Horch, K. W., 1966. Mechanism of polarized light perception, Science, Vol. 154, pp. 467-475.
- 57. Lehninger, A. L., 1959. Respiratory energy transformation, Rev. Mod. Phys., Vol. 31, pp. 136-146.
- 58. Green, D. E. and Hatefi, Y., 1961. The mitochondrion and biochemical machines, Science, Vol. 133, p. 13.
- 59. Green, D. E. and Fleischer, S., 1962. In Horizons in Biochemistry, (M. Kasha and B. Pullman, Eds.) New York and London, Adademic Press, p. 381.
- 60. Parsons, D. F., 1963. Negative staining of thinly spread cells and associated virus, J. Cell Biol., Vol. 16, pp. 620-626.
- 61. Picheral, M. B., 1966. Sur l'ultrastructure des mitochondries des cellules du tissu glandulaire du testicule de <u>Pleurodeles Waltlii Michah</u>. (Amphibien Urodele), C. R. Acad. Sc., Paris, Vol. 262, pp. 1769-1772.
- 62. Moor, H. and Muhlethaler, K., 1963. Fine structure in frozen-etched yeast cells, J. Cell Biol., Vol. 17, p. 609.

MEMBRANE ULTRASTRUCTURE IN NERVE CELLS - H. Fernández-Morán REFERENCES

- 63. Sjöstrand, F.S., Anderson-Cedergren, E. and Karlson, U., 1965. Myelin like figures formed from mitochondrial material, Nature, Vol. 202 p. 1075.
- 64. Chance, B. and Mela, L., 1966. Intramitochondrial Ph changes in cation accumulation, Proc. Natl. Acad. Sci., Vol. 55, pp. 1243-1251.
- 65. Sjöstrand, F.S. In Conference on Biological Membranes:

 Recent Progress, Section of Biological and Medical Sciences,

 New York Academy of Sciences, October, 1965, New York, in press.
- 66. André, Jean, 1965. Quelques données récentes sur la structure et la physiologie des mitochondries: glycogène, particules élémentaires, acides nucléiques, Arch. Biol. (Liège), Vol. 76, pp. 277-304.
- 67 a. Koike, M., Reed, L. and Carroll, W., 1960. J. Biol. Chem., Vol. 235, p. 1924.
 - b. Koike, M., Reed, L. and Carroll, W., 1960. J. Biol. Chem., Vol. 235, p. 1931.
- 68. Koike, M., Reed, L. and Carroll, W., 1963. J. Biol. Chem., Vol. 238, p. 30.
- 69. Fernández-Moran, H., 1964. New approaches in correlative studies of biological ultrastructure by high resolution electron microscopy, paper presented at the Celebration of the Tercentenary of the Microscope in Living Biology, the Royal Microscopical Society, Bethesda, Md., April 7-9, 1963. Published in J. Roy. Microscop. Soc., Vol. 83, Parts 1 & 2, pp. 183-195.
- 70. Fernandez-Moran, H., Reed, L., Koike, M. and Willms, C., 1964. Correlated electron microscopic and biochemical studies of a multienzyme complex: pyruvate dehydrogenase complex of <u>Escherichia coli</u>, Science, Vol. 145, pp. 930-932.
- 71. Ishikawa, C., Oliver, R. and Reed, L., 1966. α-keto acid dehydrogenase complexes, V. macromolecular organization of puruvate and α-ketoglutarate dehydrogenase complexes isolated from beef kidney mitochondria, Proc. Natl. Acac. Sci., Vol. 56, pp. 534-541.

MEMBRANE ULTRASTRUCTURE IN NERVE CELLS - H. Fernandez-Moran

REFERENCES

- 72. Fernandez-Moran, H., 1964-1965. Unpublished studies.
- 73. Caspar, D. L., and Klug, A., 1962. Physical properties in the construction of regular viruses, Cold Spring Harbor Sympos. Quant. Biol., Vol. 27, p. 1.
- 74. Tomkins, G., Yielding, K., Talal, N., and Curron, J., 1963.

 Protein structure and biological regulation, Cold

 Spring Harbor Sympos. Quant. Biol., Vol. 28, p. 461.
- 75. Hall, C. B., 1960. Measurement of globular protein molecules by electron microscopy, J. Biophys. Biochem. Cytol., Vol. 7, p. 613.
- 76. Valentine, R., 1959. The shape of protein molecules suggested by electron microscopy, Nature, Vol. 184, pp. 1838-1841.
- 77. Horne, R. W., and Greville, G. D., 1963. Observations on Ox-liver L-glutamate dehydrogenase with electron microscopy, J. Mol. Biol., Vol. 6, pp. 506-509.
- 78. Fernández-Morán, H., and Ohtsuki, M., 1966. Unpublished observations.
- 79. Haselkorn, R., Fernández-Morán, H., Kieras, F. J., and van Bruggen, E. F. J., 1965. Electron microscopic and biochemical characterization of Fraction I Protein, Science, Vol. 150, p. 1598-1601.

MEMBRANE ULTRASTRUCTURE IN NERVE CELLS - H. Fernández-Morán REFERENCES

- 80. Robertson, J. D., 1964. The Synapse: morphological and chemical correlates of function, report of NRP work session held Nov. 16-17, 1964, in Neurosciences Research Symposium Summaries (F.O. Schmitt and T. Melnechuk, eds.), Cambridge, Mass., The MIT Press, pp. 463-541.
- 81. Gray, E. G., and Guillery, R. W., 1966. Synaptic morphology in the normal and degenerating nervous system, Intn'l. Rev. Cytol., Vol. 19, pp. 111-182.
- 82. Fernández-Moran, H., 1952. The submicroscopic organization of vertebrate nerve fibres as revealed by electron microscopy. Dissertation. University of Stockholm Almquist & Wiksells AB. Uppsala.
- 83. Fernandez-Morán, H., 1952. Application of the ultrathin freezing-sectioning technique to the study of cell structures with the electron microscope, Arkiv for Fysik, Kgl. Svenska Vetenskapasakademien, Band 4, pp. 471-483.
- 84. Fernandez-Morán, H., 1953. A diamond knife for ultrathin sectioning, Exptl. Cell Res., Vol. 5, pp. 255-256.
- 85. Fernandez-Moran, H., 1956. Applications of a diamond knife for ultrathin sectioning to the study of the fine structure of biological tissues and metals, J. Biophys. Biochem. Cytol. Vol. 4, Suppl. 2, p. 29.
- 86. Fernández-Morán, H., 1960. Low temperature preparation techniques for electron microscopy of biological specimens based on rapid freezing with liquid helium II, in Annals of the New York Academy of Sciences, Vol. 85, pp. 689-713.
- 87. Fernández-Morán, H., 1960. Low temperature electron microscopy of hydrated systems, in Fast Fundamental Transfer Processes in Aqueous Biomolecular Systems (F. O. Schmitt, ed.) Cambridge, Mass., The MIT Press, pp. 33-37.
- 88. Fernandez-Morán, H., 1962. New Approaches in the Study of Biological ultrastructure by high resolution electron microscopy, paper presented at Symposium of the International Society for Cell Biology, Berne (September 1961).

MEMBRANE ULTRASTRUCTURE IN NERVE CELLS - H. Fernández-Moran REFERENCES

- 88 cont'd. <u>In</u> Symposia of the International Society for Cell Biology (R.J.C. Harris, ed.), London, Academic Press Ltd., Vol. 1, pp. 411-427.
- 89. Szent-Gyorgyi, A., 1957. Bioenergetics. New York, N. Y., Academic Press.
- 90. Frank, H. S., 1958. Covalency in the hydrogen bond and the properties of water and ice, Proc. Roy. Soc., Vol. A 247, pp. 481-492.
- 91. Hechter, O., 1964. On the role of water structures in the molecular organization of nerve cell membranes.

 Presented at American Cancer Society Conference on Cryobiology, Rye, New York, Oct. 9-10, 1964.
- 92. Fernández-Morán, H., 1961. High resolution electron microscopy of hydrated biological systems, Proc. Internat. Biophys. Congr., Stockholm, Sweden.
- 93. Fernández-Morán, H., 1960. Direct study of ice crystals and of hydrated systems by low-temperature electron microscopy, J. Appl. Phys., Vol. 31, p. 1841.
- 94. Klotz, I. M., 1960. Protein molecules in solution, Circulation, Vol. 21, p. 828.
- 95. Pauling, L., 1961. A molecular theory of general anesthesia, Science, Vol. 134, p. 15.
- 96. Bernard, C., 1875. Leçons sur les anesthesiques et sur l'asphyxie, Paris, Bailliere, France.
- 97. Egelstaff, P. A., 1962. Advances Phys., Vol. 11, p. 203.
- 98. Fernández-Morán, H., 1965. Forms of water in biologic systems and the organization of membranes, in Annals of the New York Academy of Sciences, Vol. 125, Art. 2, p. 739-753.
- 99. Berendsen, H. J. C., and Migchelson, 1965. Hydration structure of fibrous macromolecules, in Annals of the New York Academy of Sciences, Vol. 125, Art. 2, pp. 365-379.
- 100. Eigen, M., and DeMayer, L., 1959. Hydrogen bond structure, proton hydration, and proton transfer in aqueous solution.

 <u>In</u> the Structure of Electrolytic Solutions (W. J. Hammer, ed.), New York, N. Y., John Wiley & Sons, Inc., pp. 64-85.

MEMBRANE ULTRASTRUCTURE IN NERVE CELLS - H. Fernández-Morán REFERENCES

- 101. Swift, H., Kislev, N., and Bogorad, L., 1964. Evidence for DNA and RNA in mitochondria and chloroplasts, J. Cell Biol., Vol. 23, 91A.
- 102. André, J., and Marinozzi, V., 1965. Presence, dans les mitochondries, de particules ressemblant aux ribosomes, J. Microscopie, Vol. 4, pp. 615-626.
- 103. Van Bruggen, E.F.J., Borst, P., Ruttenberg, G.J.C.M., Gruber, M., and Kroon, A.M., 1966. Circular mitochondrial DNA, Biochem. Biophys.Acta, Vol. 119, p. 437.
- 104. Nass, M.M.K., 1966. The circularity of mitochondrial DNA, Proc. Nat'l. Acad. Sci., Vol. 56, pp. 1215-1222.
- 105. Colvill, A. J., Van Bruggen, E.F.J., and Fernandez-Moran, H., 1966. Physical properties of a DNA-dependent RNA polymerase from Escherichia coli, J. Mol. Biol., Vol. 17, pp. 302-304.
- 106. Slayter, H., and Hall, C., 1966. Electron microscopy of RNA polymerase and RNA polymerase bound to T7 DNA, J. Mol. Biol., Vol. 21, pp. 113-114.
- 107. a. Richardson, John, 1966. The binding of RNA polymerase to DNA, J. Mol. Biol., Vol. 21, pp. 83-114.
 - b. Richardson, John, 1966. Enzymic synthesis of RNA from T7 DNA, ibid, pp. 115-127.
- 108. Schmitt, F. O., 1964. Molecular and Ultrastructural correlates of function in neurons, neuronal nets, and the brain, discussion paper in Neurosciences Research Symposium Summaries (F.O. Schmitt and T. Melnechuk, eds.), Cambridge, Mass., The MIT Press, pp. 321-351.
- 109. Lehninger, A. L., 1963. Cell membranes, a report of an NRP Work Session, July 27-28, 1963, in Neurosciences Research Symposium Summaries (F.O. Schmitt and T. Melnechuk, eds.), Cambridge, Mass., the MIT Press, pp. 1-57.
- 110. Lehninger, A. L., 1964. Supramolecular Organization of Enzyme and membrane systems, paper presented at NRP work session, September 29, 1964, in Neurosciences Research Symposium Summaries (F.O. Schmitt and T. Melnechuk, eds.) Cambridge, Mass., The MIT Press, pp. 294-317.

MEMBRANE ULTRASTRUCTURE IN NERVE CELLS - H. Fernández-Morán REFERENCES

- 111. Kavanau, J. L., 1965. Structure and Function in Biological Membranes, San Francisco, Holden-Day.
- 112. Eigen, M., and DeMayer, L.C.M., 1965. Information storage and processing in biomolecular systems, summary of two NRP work sessions held May 5, 1965, in Neurosciences Research Symposium Summaries (F.O. Schmitt, and T. Melnechuk, eds.), Cambridge, Mass., The MIT Press, pp. 244-266.
- 113. Fernandez-Moran, H., 1965. Electron microscopy with high field superconducting solenoid lenses, Proc. Nat'l. Acad. Sci., Vol. 53, pp. 445-451.
- 114. Fernandez-Morán, H., 1966. High-resolution electron microscopy with superconducting lenses at liquid helium temperatures, Proc. Nat'l. Acad. Sci., Vol. 56, pp. 801-808.
- 115. Dupouy, G., and Perrier, F., 1966. Amelioration du contraste en microscopie electronique, Proc. Sixth Intn'l. Cong. for Electron Microscopy, Kyoto, Japan, Aug. 28-Sept. 4, 1966, Vol. I, p. 3.
- 116 a. Fernandez-Morán, H., 1966. High resolution electron microscopy of biological specimens, Proc. Sixth Intn'l. Cong. for Electron Microscopy, Kyoto, Japan, Aug. 28-Sept. 4, 1966, Vol. I, p. 13.
 - b. Fernández-Morán, H., 1966. Applications of Improved point cathode sources to high resolution electron microscopy, ibid, p. 27.
 - c. Fernandez-Morán, H., 1966. Low temperature electron microscopy with high field superconducting lenses, <u>ibid</u>, p. 147.
- 117. Möllenstedt, G., and Speidel, R., 1960. Elektronenoptischer mikroschreiber unter elektronenmikroskopischer arbeitskontrolle, Phys. Blät., Vol. 16, pp. 192-198.
- 118. Fernandez-Morán, H., 1966. Potential application of electron optical methods to storage of information for direct retrieval, in <u>Biology and the Exploration of Mars</u>,

REFERENCES

- 118. cont'd. National Academy of Sciences--National Research Council, Publication No. 1296, Washington, D.C.
- 119 a. Calvin, M., 1959. Energy reception and transfer in photosynthesis, Revs. Mod. Phys., Vol. 31, p. 147.
 - b. Calvin, M., 1959. Free radicals in photosynthetic systems, ibid, pp. 157-161.
- 120. Arnold, W. and Clayton, R.K., 1960. The first step in photosynthesis: evidence for its electronic nature, Proc. Nat'l. Acad. Sci., Vol 46, p. 769.
- 121. Parson, P. and Simpson, M.V., 1967. Biosynthesis of DNA by isolated Mitochondria: Incorporation of Thymidine Triphosphate-2-C¹⁴. Science, 155, pp. 91-93.
- 122. Kellenberger, E., 1966. Control Mechanisms in Bacteriophage Morphopoiesis in "Principles of Biomolecular Organization" (Wolstenholme, G.E.W. and O'Connor, M. editors) J.A. Churchill, Ltd.
- 123. Katchalsky, A. and Oplatka, A., 1966. Hysteresis and Macromolecular Memory in "Neurosciences Research Symposium Summaries" (Schmitt, F.O. and Melnechuck, T. editors) M.I.T. press, Vol. 1, pp. 352-374.
- 124 a. von Neumann, J., 1956. Probabilistic Logics and the Synthesis of Reliable Organisms from Unreliable Components, Automata Studies, Annals of Mathematics Studies, Study 34, Princeton Univ. Press, pp. 43-98.
 - b. von Neumann, J., 1966. Theory of Self-Reproducing Automata (Burks, A.W., edited and completed), Univ. of Ill. Press, Urbana, Ill. and London.
- 125. Monod, J., 1966. From Enzymatic Adaption to Allosteric Transitions, Science, 154, pp. 475-483.
- 126. Luse, S.A., 1962. Ultrastructure of the Brain and its Relation to Transport of Metabolites in "Ultrastructure and Metabolism of the Nervous System", Proc. of Assoc. for Research in Nervous and Mental Disease, The Williams and Wilkins Co., Vol. XL, pp. 1-26.

MEMBRANE ULTRASTRUCTURE IN NERVE CELLS - H. Fernández-Morán REFERENCES

- 127. Nachmansohn, D., 1959. Chemical and Molecular Basis of Nerve Activity, Academic Press, N.Y.
- 128. Shanthaveerappa, T.R. and Bourne, G.H., 1966. Perineural Epithelium: A New Concept of its Role in the Integrity of the Peripheral Nervous System, Science, 154, pp. 1464-1467.
- 129. Hyden, H., personal communication.
- 130. Crain, S.M. and Peterson, E.R., 1964. Complex bioelectric activity in organized tissue cultures of spinal cord (human, rat and chick). J. Cell. Comp. Physiol. 64, pp. 1-13.
- 131. Pease, R.F.W., Hayes, T.L., Camp, A.S. and Amer, N.M., 1966. Electron Microscopy of Living Insects, Science, <u>154</u>, pp. 1185-1186.
- 132. Changeux, J.P, Thiery, J., Tung, Y. and Kittel, C., 1966.
 On the Cooperativity of Biological Membranes. Proc.
 Nat'l. Acad. Sci., Vol. 56, in press.
- 133. Anderson, N.G., 1966. The development of zonal centrifuges,
 Journal of the National Cancer Institute, Monograph No. 21
 (U.S.A.).
- 134. Fernandez-Morán, H., 1951. Las Bases Físicas y Fisiológicas del Electroencefalograma. Anais do IV Congresso Sul-Americano de Neuro-cirugía, Porto Alegre, Brasil, pp. 37; and Acta Cientifica Venezolana, 2, pp. 94-102.
- 135. Polonsky, J., 1962. On the Origin of Information and Organization in Polyatomic Structures. Annales de Radioelectricite, XVII, pp. 227-245.




FIG.1 (a) High-resolution electron micrograph of myelin sheath segment from transverse thin section of frog sciatic nerve showing concentric array of dense and intermediate layers. Particulate subunit structures (arrows) are regularly found within the plane of the layers in these well-preserved, osmium fixed, low-temperature preparations. X 480,000. (b) Axonal mitochondria from a similar preparation demonstrating globular subunits in the dense layers of the cristae. X 600,000. (c) Low-angle x-ray diffraction pattern of fresh rat sciatic nerve featuring a fundamental period of 178Å, with typical alternation of the intensities of the even and odd orders.




FIG.2 (a) High-resolution micrograph of thin myelin sheath segment from transverse section of frog sciatic nerve, fixed and embedded at low temperatures. The characteristic particulate fine structure (arrows) detected mainly in the dense intermediate layers is also revealed by other techniques yielding improved preservation of the highly regular laminated sheath structure. X 900,000. (b,c) Ultrathin frozen section of fresh frog sciatic nerve negatively stained with phosphotungstate and examined by low temperature electron microscopy. These repeating particulate units (arrows) attached to the dense myelin layers have consistently been found only in negatively stained, unfixed, specimens. Bearing in mind the inherent artifact possibilities, their possible relationship to the subunit structures, already shown by other methods, is now being further investigated. X 1,300,000.




FIG. 3 High-resolution electron micrographs of thin sections of frog retinal rod outer segments: (a) Osmium-fixed, low-temperature preparation of dark-adapted retina showing arrays of dense particles about 40Å in diameter (arrows) within the compound membrane of the unit discs, X700,000; (b,c) ultrathin sections of fresh, unfixed, light-adapted rod outer segments prepared without embedding by improved techniques, and stained with uranyl formate. The ordered spherical particles with indications of substructure of 10-15Å (arrows) can be detected in the dense and intermediate membrane layers. The presence of an ordered substructure within these membranes is substantiated by recent x-ray diffraction studies. X 900,000.




FIG. 4. Electron micrographs of: (a)enlarged segment of crista from negatively stained beef heart mitochondrion demonstrating three parts of the elementary particle (EP): head piece, stalk, and base piece, X 900,000; (b) segment of crista from thin mitochondrial section showing arrays of similar particles, X 700,000; (c)particulate units of approximately the same size as the EP, containing the electron transfer chain, isolated from beef heart mitochondria by Dr. D.E. Green et al., X 2,000,000. (d)Diagrammatic illustration of a macromolecular assembly containing many species of macromolecular components. Courtesy of Professor F.O. Schmitt.




FIG. 5. High resolution electron micrographs of: (a) E. coli pyruvate dehydrogenase complex (PDC) prepared by $\overline{\text{Dr. L.J.}}$ Reed et al. showing characteristic central $\overline{\text{tetrad}}$ and associated subunits of this multienzyme complex negatively stained with phosphotungstate, and (b) positively stained with uranyl acetate, X 1,300,000. (c,d) E. coli α -ketoglutarate dehydrogenase complex (KGDC) isolated in Dr. Reed's laboratory displaying similar structural features of the positively stained particles in different orientations, X 1,400,000.




FIG. 6. Electron micrographs of bovine L-glutamate dehydrogenase (GDH) (Boehringer & Soehne, Mannheim): (a) microcrystals negatively stained with uranyl formate by microdroplet cross-spraying technique showing typical periodic lattice formed through ordered aggregation of the constituent molecules, X 600,000; (b) in thinner, ribbon-like strands the regular substructure of individual molecules (arrow) can be discerned, X 600,000; (c,d) as a result of the improved preservation and enhanced resolution achieved in these preparations, undissociated individual molecules (ca. 1 million molecular weight) are frequently detected, appearing as spherical or polyhedral particles of diameter 130 - 150 Å in different orientations. Subunits of triangular shape and size are also found. X 900,000. In addition to its specific value investigation of this enzyme is also of interest as a model system to illustrate the different states of molecular and supramolecular association bearing on the general organization of enzyme complexes in membranes




FIG. 7. (a) Electron micrograph of shadow-cast particles of Fraction I protein isolated from Chinese cabbage leaves by F.J. Kieras and R. Haselkorn. This major soluble protein fraction located predominantly in the chloroplasts has carboxydismutase activity. As seen in negatively stained preparations (b) the individual particle appears to be a cube, about 120Å along each edge, containing 24 subunits. This is one of the first demonstrations that the substructure of a protein enzyme can be directly resolved in the size range of the enzyme complexes associated with membrane systems.




FIG. 8. Electron micrographs of: (a) fine submicroscopic nerve fiber from the lateral funiculus of the frog spinal cord. The fiber shaft has a width of 1000 Å and consists of a flattened, ultrathin tubular membrane (NM) enclosing an individual axon filament (Af) 200 Å in diameter. X 54,000; (b) Submicroscopic nerve fiber ending and associated membrane components isolated from fresh frog spinal cord, stained with uranyl formate, and examined by low-temperature electron microscopy. These well-preserved ultrathin nerve membrane preparations exhibit a characteristic surface structure featuring in certain regions hexagonal arrays of closely packed polygonal elements, 60 to 90 Å in diameter (insert). Various types of larger particulate components (arrows) are found attached to the membranes, particularly in junctional regions. X 100,000. Insert: X 600,000.




FIG. 9. High resolution electron micrographs of: (a) submicroscopic nerve fiber ending isolated from fresh frog spinal cord using improved preparation techniques which permit examination of whole mounts of ultrathin nerve membranes, without introducing the fixation, extraction, and embedding artifacts inherent in standard thin-sectioning methods. Under these favorable conditions an underlying subunit structure of the membrane surfaces can be resolved, with associated electron-dense components (arrows), X 400,000; (b) as shown in the enlarged segment of this membrane, numerous polygonal components with electron-dense or annular cores (arrows), 100 to 200 Å in diameter, contribute to the mosaic-like surface pattern, X 800,000. Further methodological improvements should eventually make it possible to directly observe the molecular organization of individual nerve membranes in their native hydrated state.




FIG. 10. Low-temperature electron micrographs of: (a) lecithin micelles embedded in thin, buffered phosphotungstate film showing periodic dense lines separated by light bands corresponding to the hydrophilic and hydrophobic regions of adjacent bimolecular leaflets; (b) lecithin micelles similarly embedded in thin phosphotungstate film, and exposed to argon gas under pressure, showing dense particulate aggregates (arrows) along the dense and intermediate layers, which may represent the resulting argon hydrate microcrystals in the size range of 10 -20 Å. X 2,000,000. Model experiments of this kind could provide direct experimental evidence for the existence of hydrate microcrystals forming reversibly in nerve membranes as first postulated by L. Pauling.



FIG. 11. Electron micrograph of thin section through retinulae from a tropical "Skipper" butterfly demonstrating the highly regular organization of the differentiated membrane-bound components in these photoreceptors. The matched rhabdomere pairs (arrows-R) are surrounded by rows of mitochondria(Mi), grouped together in radial columns by a network of fine tubular channels(Ut) arising from the tracheal compartments (T). The walls of the tubular rhabdomere compartments (insert) contain arrays of dense particles which may correspond to the ordered arrangement of the dichroic photopigment molecules providing the basis for the polarization-analyzing property of the insect eye.




FIG. 12. Electron micrographs showing: (a) characteristic attachment of RNA polymerase molecules (arrows) to DNA strands as seen in shadowed preparations, X 400,000; (b) DNA-dependent RNA polymerase molecules from E. coli prepared by the Chamberlin & Berg procedure appear to consist in negatively stained preparations of six subunits arranged in a hexagon with a cross-section of 120 to 130 Å. These correlated electron microscopic and biochemical studies are expected to yield further information on RNA polymerase and its participation in the differential RNA transcription upon DNA templates, which is of fundamental importance in the regulation of protein synthesis and function in nerve cell membranes and their derivatives.




FIG. 13. High resolution electron micrographs showing: (a) the complex arrangement of the structural subunits of about 8 - 20 Å within the periodic lattice of catalase crystals stained with uranyl formate, X 1,200,000; (b) phase contrast image of the hexagonal array of carbon atoms in ultrathin layer of single-crystal graphite, X 32,000,000. This level of resolution of pauciatomic structural patterns, which can now be achieved under certain conditions with improved instrumentation and preparation techniques, should prove to be of key value in correlated studies for direct visualization of the molecular organization of nerve membranes and associated enzyme systems.

FIGURE CAPTIONS

- Fig. 1. (a) High resolution electron micrograph of myelin sheath segment from transverse thin section of frog sciatic nerve showing concentric array of dense and intermediate layers. Particulate sub-unit structures (arrows) are regularly found within the plane of the layers in these well-preserved, osmium fixed, low-temperature preparations. X 320,000. (b) Axonal mitochondria from a similar preparation demonstrating globular subunits in the dense layers of the cristae. X 400,000. (c) Low-angle x-ray diffraction pattern of fresh rat sciatic nerve featuring a fundamental period of 178 Å, with typical alternation of the intensities of the even and odd orders.
- Fig. 2. (a) High resolution micrograph of thin myelin sheath segment from transverse section of frog sciatic nerve, fixed and embedded at low temperatures. The characteristic particulate fine structure (arrows) detected mainly in the dense intermediate layers is also revealed by other techniques yielding improved preservation of the highly regular laminated sheath structure. X 600,000. (b,c) Ultrathin frozen section of fresh frog sciatic nerve negatively stained with phosphotungstate and examined by low temperature electron microscopy. These repeating

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Morań particulate units (arrows) attached to the dense myelin layers have consistently been found only in negatively stained, unfixed, specimens. Bearing in mind the inherent artifact possibilities, their possible relationship to the subunit structures, already shown by other methods, is now being further investigated. X 870,000.

Fig. 3. High resolution electron micrographs of thin sections of frog retinal rod outer segments: (a) Osmium-fixed, low-temperature preparation of dark-adapted retina showing arrays of dense particles about 40 Å in diameter (arrows) within the compound membrane of the unit discs. X 470,000. (b,c) Ultrathin sections of fresh, unfixed, light-adapted rod outer segments prepared without embedding by improved techniques, and stained with uranyl formate. The ordered spherical particles with indications of substructure of 10-15 Å (arrows) can be detected in the dense and intermediate membrane layers. The presence of an ordered substructure within these membranes is substantiated by recent x-ray diffration studies. X 600,000.

- Fig. 4. Electron migrographs of: (a) enlarged segment of crista from negatively stained beef heart mito-chondrion demonstrating three parts of the elementary particle (EP): head piece, stalk, and base piece, X600,000; (b) segment of crista from thin mitochondrial section showing arrays of similar particles, X470,000; (c) particulate units of approximately the same size as the EP, containing the electron transfer chain, isolated from beef heart mitochondria by Dr. D.E. Green et al., X1,330,000; (d) diagrammatic illustration of a macromolecular assembly containing many species of macromolecular components. Courtesy of Professor F. O. Schmitt.
- Fig. 5. High resolution electron micrograph of: (a) E. coli pyruvate dehydrogenase complex (PDC) prepared by

 Dr. L. J. Reed et al. showing characteristic central tetrad and associated subunits of this multienzyme complex negatively stained with phosphotungstate, and (b) positively stained with uranyl acetate,

 X 870,000. (c,d) E. coli α-ketoglutarate dehydrogenase complex (KGDC) isolated in Dr. Reed's laboratory displaying similar structural features of the positively stained particles in different orientations, X 930,000.

- Fig. 6. Electron micrographs of bovine L-glutamate dehydrogenase (GDH) (Boehringer & Soehne, Mannheim): (a) microcrystals negatively stained with uranyl formate by microdroplet cross-spraying technique showing typical periodic lattice formed through ordered aggregation of the constituent molecules, X 400,000; (b) in thinner, ribbon-like strands the regular substructure of individual molecules (arrow) can be discerned, X 400,000; (c,d) as a result of the improved preservation and enhanced resolution achieved in these preparations, undissociated individul molecules (ca. 1 million molecular weight) are frequently detected, appearing as spherical or polyhedral particles of diameter 130 - 150 Å in different orientations. Subunits of triangular shape and size are also found. X 600,000. In addition to its specific value investigation of this enzyme is also of interest as a model system to illustrate the different states of molecular and supramolecular association bearing on the general organization of enzyme complexes in membranes.
- Fig. 7. (a) Electron micrograph of shadow-cast particles of Fraction I protein isolated from Chinese cabbage leaves by F. J. Kieras and R. Haselkorn. This major soluble protein fraction located predominantly in the chloroplasts has carboxydismutase activity. X 120,000.

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Morán

As seen in negatively stained preparations (b) the individual particle appears to be a cube, about 120 Å along each edge, containing 24 subunits. X 400,000. This is one of the first demonstrations that the substructure of a protein enzyme can be directly resolved in the size range of the enzyme complexes associated with membrane systems.

Fig. 8. Electron micrographs of: (a) fine submicroscopic nerve fiber from the lateral funiculus of the frog spinal cord. The fiber shaft has a width of 1000 A and consists of a flattened, ultrathin tubular membrane (NM) enclosing an individual axon filament (Af) 200 A in diameter. X 36,000; (b) submicroscopic nerve fiber ending and associated membrane components isolated from fresh frog spinal cord, stained with uranyl formate, and examined by low-temperature electron microscopy. These well-preserved ultrathin nerve membrane preparations exhibit a characteristic surface structure featuring in certain regions hexagonal arrays of closely packed polygonal elements, 60 to 90 A in diameter (insert). Various types of larger particulate components (arrows) are found attached to the membranes, particularly in junctional regions. X 670,000. Insert: X 400,000.

- High resolution electron micrographs of: (a) sub-Fig. 9. microscopic nerve fiber ending isolated from fresh frog spinal cord using improved techniques which permit examination of whole mounts of ultrathin nerve membranes, without introducing the fixation, extraction, and embedding artifacts inherent in standard thin-sectioning methods. Under these favorable conditions an underlying subunit structure of the membrane surface can be resolved, with associated electron-dense components (arrows), X 270,000; (b) as shown in the enlarged segment of this membrane, numerous polygonal components with electron-dense or annular cores (arrows), 100 to 200 A in diameter, contribute to the mosaic-like surface pattern, X 530,000. Further methodological improvements should eventually make it possible to directly observe the molecular organization of individual nerve membranes in their native hydrated state.
- Fig. 10. Low-temperature electron micrographs of: (a)

 lecithin micelles embedded in thin, buffered

 phosphotungstate film showing periodic dense lines

 separated by light bands corresponding to the hydro
 philic and hydrophobic regions of adjacent bio
 molecular leaflets; (b) lecithin micelles similarly

Membrane Ultrastructure in Nerve Cells - H. Fernandez-Moran

embedded in thin phosphotungstate film, and exposed

to argon gas under pressure, showing dense particulate
aggregates (arrows) along the dense and intermediate
layers, which may represent the resulting argon
hydrate microcrystals in the size range of 10 - 20 Å.

X 1,330,000. Model experiments of this kind could
provide direct experimental evidence for the existence
of hydrate microcrystals forming reversibly in nerve

membranes as first postulated by L. Pauling.

Fig. 11. Electron micrograph of thin section through retinulae from a tropical "Skipper" butterfly demonstrating the highly regular organization of the differentiated membrane-bound components in these photoreceptors.

The matched rhabdomere pairs (arrows-R) are surrounded by rows of mitochondria (Mi), grouped together in radial columns by a network of fine tubular channels (Ut) arising from the tracheal compartments (T). The walls of the tubular rhabdomere compartments (insert) contain arrays of dense particles which may correspond to the ordered arrangement of the dichroic photopigment molecules providing the basis for the polarization-analyzing property of the insect eye. X 12,390. Insert: X 354,000.

- Fig. 12. Electron micrographs showing: (a) characteristic attachment of RNA polymerase molecules (arrows) to DNA strands as seen in shadowed preparations, X 284,000; (b) DNA-dependent RNA polymerase molecules from E. coli prepared by the Chamberlin & Berg procedure appear to consist in negatively stained preparations of six subunits arranged in a hexagon with a cross-section of 120 to 130 Å. X 2,130.000. These correlated electron microscopic and biochemical studies are expected to yield further information on RNA polymerase and its participation in the differential RNA transcription upon DNA templates, which is of fundamental importance in the regulation of protein synthesis and function in nerve cell membranes and their derivatives.
- Fig. 13. High resolution electron micrographs showing: (a)

 the complex arrangement of the structural subunits

 of about 8 20 Å within the periodic lattice of

 catalase chystals stained with uranyl formate,

 X 760,000; (b) phase contrast image of the hexagonal

 array of carbon atoms in ultrathin layer of single
 crystal graphite, X 20,290,000. This level of

 resolution of pauciatomic structural patterns, which

 can now be achieved under certain conditions with

 improved instrumentation and preparation techniques,

 should prove to be of key value in correlated studies

 for direct visualization of the molecular organization

 of nerve membranes and associated enzyme systems.

ACKNOWLEDGEMENTS

I am much indebted to Miss J. E. Richardson for valuable help in editing and preparing the manuscript, and to Messrs. C. L. Hough and C. Weber for expert assistance with all photographic reproductions. Sincere thanks are also due to Messrs. M. Ohtsuki and R. Vicario, and to Misses D. A. Meddoff, V. Manrique, A. Gustafson, and Mrs. S. Erikson for their skillful assistance in the course of this work.

It is a pleasure to thank Prof. F. O. Schmitt of the Massachusetts Institute of Technology, Prof. W. H. Sweet of the Massachusetts General Hospital, Prof. H. Hyden of the University of Göteborg, Sweden, Prof. A. Katchalsky of the Weizmann Institute of Science, Israel, Prof. H. S. Bennett of the University of Chicago, and Dr. O. Hechter of the Worcester Foundation for Experimental Biology, for many valuable suggestions and stimulating discussions.

^{*} This work was supported by grant GM-13243 from the National Institutes of Health, by grant NsG 441-63 of the National Aeronautics and Space Administration, by United States Atomic Energy Commission contract AT (11-1) - 1344, by the L. Block Fund, and by the Otho Sprague Memorial Fund of the University of Chicago.